


[User Agreement - Privacy Policy](#)

**Support the
Mental Health
Crisis Response
Fund!**

[Email Alert Sign-up!](#)

Leading The Way For America's Mental Health

Go!

Post-Traumatic Stress Disorder


Post-traumatic stress disorder (PTSD) is an extremely debilitating condition that can occur after exposure to a terrifying event or ordeal in which grave physical harm occurred or was threatened. Traumatic events that can trigger PTSD include violent personal assaults such as rape or mugging, natural or human-caused disasters, accidents, or military combat.

Military troops who served in Vietnam and the Gulf Wars; rescue workers involved in the aftermath of the Oklahoma City Bombing; survivors of accidents, rape, physical and sexual abuse, as well as other crimes; immigrants fleeing violence in their home countries; survivors of the 1994 California earthquake, the 1997 South Dakota floods, and hurricanes Hugo and Andrew; and people who witness traumatic events are among the people who may develop PTSD. Families of victims can also develop the disorder.

Fortunately, through research supported by the National Institute of Mental Health (NIMH) and the Department of Veterans Affairs (VA), effective treatments have been developed to help people with PTSD. Research is also helping scientists better understand the condition and how it affects the brain and the rest of the body.

What Are the Symptoms of PTSD?

Many people with PTSD repeatedly re-experience the ordeal in the form of flashback episodes, memories, nightmares, or frightening thoughts, especially when they are exposed to



**Confidential
depression-
screening test**

- [Time for Reassurance](#)
- [When to Seek Help](#)
- [Helping Children Handle Disaster-Related Anxiety](#)
- [Helping Children Cope with Loss](#)
- [Young Adults](#)
- [College Students](#)
- [Adults](#)
- [Older Adults](#)
- [Coping with Loss](#)
- [Post-Traumatic Stress Disorder](#)
- [Helping Your Workforce Cope and Return to Work](#)
- [People of Faith](#)

events or objects reminiscent of the trauma. Anniversaries of the event can also trigger symptoms. People with PTSD also experience emotional numbness and sleep disturbances, depression, anxiety, and irritability or outbursts of anger. Feelings of intense guilt are also common. Most people with PTSD try to avoid any reminders or thoughts of the ordeal. PTSD is diagnosed when symptoms last more than one month.

How Common Is PTSD?

At least 3.6% of U.S. adults (5.2 million Americans) have PTSD during the course of a year. About 30 percent of the men and women who have spent time in war zones experience PTSD. One million war veterans developed PTSD after serving in Vietnam. PTSD has also been detected among veterans of the Persian Gulf War, with some estimates running as high as 8 percent.

When Does PTSD First Occur?

PTSD can develop at any age, including in childhood. Symptoms typically begin within 3 months of a traumatic event, although occasionally they do not begin until years later. Once PTSD occurs, the severity and duration of the illness varies. Some people recover within 6 months, while others suffer much longer.

- [Tips for Primary Care Physicians - Talking with your Patients about Trauma](#)
- [Newsreleases](#)

What Treatments Are Available for PTSD?

Research has demonstrated the effectiveness of cognitive-behavioral therapy, group therapy, and exposure therapy, in which the patient repeatedly relives the frightening experience under controlled conditions to help him or her work through the trauma, as well as medications that help ease the symptoms of depression and anxiety and help promote

sleep. Scientists are attempting to determine which treatments work best for which type of trauma.

Do Other Physical or Emotional Illnesses Tend to Accompany PTSD?

Co-occurring depression, alcohol or other substance abuse, or another anxiety disorder are not uncommon. The likelihood of treatment success is increased when these other conditions are appropriately diagnosed and treated as well.

Headaches, gastrointestinal complaints, immune system problems, dizziness, chest pain, or discomfort in other parts of the body are common. Often, doctors treat the symptoms without being aware that they stem from PTSD. NIMH, through its education program, is encouraging primary care providers to ask patients about experiences with violence, recent losses, and traumatic events, especially if symptoms keep recurring. When PTSD is diagnosed, referral to a mental health professional who has had experience treating people with the disorder is recommended.

Who Is Most Likely to Develop PTSD?

People who have been abused as children or who have had other previous traumatic experiences are more likely to develop the disorder. Research is continuing to pinpoint other factors that may lead to PTSD.

What Are Scientists Learning From Research?

NIMH and the VA sponsor a wide range of basic, clinical, and genetic studies of PTSD. In addition, NIMH has a special finding mechanism, called RAPID Grants, which allows researchers to immediately visit the scenes of disasters, such as plane crashes or floods and hurricanes, to study the acute effects of the event and the effectiveness of early intervention.

Research has shown that PTSD clearly alters a number of fundamental brain mechanisms. Because of this, abnormalities have been detected in brain chemicals that mediate coping behavior, learning, and memory among people with the disorder. Recent brain imaging studies have detected altered metabolism and blood flow as well as anatomical changes in people with PTSD.

Post-Traumatic Stress Disorder

The following are also recent research findings:

- Some studies show that debriefing people very soon after a catastrophic event may reduce some of the symptoms of PTSD. A study of 12,000 schoolchildren who lived through a hurricane in

Hawaii found that those who got counseling early on were doing much better two years later than those who did not.

- People with PTSD tend to have abnormal levels of key hormones involved in response to stress. Cortisol levels are lower than normal and epinephrine and norepinephrine are higher than normal. Scientists have also found that people with this condition have alterations in the function of the thyroid and in neurotransmitter activity involving serotonin and opiates.
- When people are in danger, they produce high levels of natural opiates, which can temporarily mask pain. Scientists have found that people with PTSD continue to produce those higher levels even after the danger has passed; this may lead to the blunted emotions associated with the condition.
- It used to be believed that people who tend to dissociate themselves from a trauma were showing a healthy response, but now some researchers suspect that people who experience dissociation may be more prone to PTSD.
- Animal studies show that the hippocampus -- a part of the brain critical to emotion-laden memories -- appears to be smaller in cases of PTSD. Brain imaging studies indicate similar findings in humans. Scientists are investigating whether this is related to short-term memory problems. Changes in the hippocampus are thought to be responsible for intrusive memories and flashbacks that occur in people with this disorder.
- Research to understand the neurotransmitter system involved in memories of emotionally charged events may lead to discovery of drugs that, if given early, could block the development of PTSD symptoms.
- Levels of CRF, or corticotropin releasing factor--the ignition switch in the human stress response--seem to be elevated in people with PTSD, which may account for the tendency to be easily startled. Because of this finding, scientists now want to determine whether drugs that reduce CRF activity are useful in treating the disorder.

The content of this fact sheet was adapted from material published by the National Institute of Mental Health.

For more information contact:

**National Mental Health
Association**

800.969.NMHA

<http://www.nmha.org>

Other Resources:

National Institute of Mental Health **888.8.ANXIETY**

<http://www.nimh.nih.gov/anxiety/>

Obsessive-Compulsive Foundation **203.878.5669**

<http://www.ocfoundation.org>

Anxiety Disorders Association of America **301.231.9350**

<http://www.adaa.org>

Freedom From Fear **718.351.1717**

<http://www.freedomfromfear.com/>

American Psychiatric Association **202.682.6000**

<http://www.psych.org>

American Psychological Association **800.964.2000**

<http://www.apa.org>

<http://helping.apa.org/>

American Academy of Child and Adolescent Psychiatry **800.333.7636
Ext. 124**

<http://www.aacap.org>

© 2001 National Mental Health Association