



UCSU 20/6.22/2.903/1998
c.2



G A R D E N I N G  S E R I E S

COLORADO STATE PUBLICATIONS LIBRARY
UCSU20/6.22/2.903/1998 c.2 local
Pottorff, Laura Pic/Nonchemical disease



3 1799 00025 1132

DISEASES

Nonchemical disease control

no. 2.903

by Laura Pickett Pottorff

Quick Facts...

Many disease problems can be prevented or controlled without the use of pesticides.

Always choose varieties of plants that are adapted to Colorado growing conditions.

Avoid bringing diseases into the garden or moving them around (exclusion).

Eliminate the disease-causing organism after it has become established on a plant.

With the increasing concern about use and misuse of pesticides in commercial agriculture and home gardens, there are more and more inquiries for "organically grown" commodities every year. Nonchemical control practices for plant diseases have been known and recommended for years. The backbone of any integrated pest control program must always include cultural and sanitation practices, two important components of nonchemical disease control.

Unfortunately, disease problems may begin as soon as seeds are planted and can continue into harvest and storage. Plant diseases may be caused by several different living pathogenic organisms such as fungi, bacteria, viruses, mycoplasmas and nematodes. In addition to these parasitic organisms, non-living factors such as deficiencies or excesses of water, light, temperatures, air pollution, pesticides and nutrients can either predispose a plant to disease or directly cause plant injury.

Fortunately, many disease problems can be prevented or controlled without the use of pesticides. Effective plant disease control must begin at the onset of disease or even before symptoms appear.

Resistance

Effective plant disease control through resistance (or a plant's tolerance or immunity to a disease) is based on the knowledge of diseases known to occur in an area. Always choose varieties of plants that are adapted to Colorado growing conditions. Many vegetables, fruit, and ornamental plant varieties are available with resistance to one or more diseases.

For example, always select plants labeled "VFN," "VFNA," "VFNT," etc., when purchasing tomato varieties. This indicates that the plants are resistant to Verticillium wilt (V), Fusarium wilt (F), Southern root-knot nematode (N), early blight (A), or tobacco (tomato) mosaic virus (T). Selection of resistant plants may eliminate many disease problems. Contact your local Colorado State University Cooperative Extension county office for lists of plant varieties that are successfully grown in this area.

Exclusion

Exclusion is preventing the entrance and establishment of disease-causing organisms (pathogens) into areas where plants are grown. This means to avoid bringing diseases into the garden or moving them around.

Use certified, disease-free seed or transplants. Examine the leaves and root systems of transplants and eliminate or destroy diseased plants. Either raise your own transplants in sterilized beds or buy them from a reputable dealer. Do not purchase transplants with galls or swellings on their roots or plants that have a brown discoloration on the stem at the ground line. Galls or swellings may

**Colorado
State**
University[®]
Cooperative
Extension

© Colorado State University
Cooperative Extension. 6/92.

indicate root-knot nematode infection and a brown stem discoloration may denote the presence of damping-off organisms.

Also, avoid transporting soil or tools from known disease areas to disease-free areas.

Eradication

Eradication is the elimination of the disease-causing organism after it has established on a plant. Eradication can be accomplished by several methods.

Sanitation

Plant pathogens are less likely to survive if organic matter is quickly decomposed. Remove plant debris or infected plant parts after each growing season. Turn the soil after harvest to help break down small roots that may harbor nematode, fungal or bacterial pathogens. Gardeners may compost dead plants if they have a good composting system; otherwise these piles may serve as an inoculum source of pathogens.

Prune or remove twigs and branches of woody plants affected with fire blight and other bacterial or fungal canker diseases. (See fact sheet 2.907, *Fire blight*.)

Keep gardens weed-free. Weeds often are another source of pathogen inoculum. Eradication of weeds can break the life cycle of a pathogen and control it. Weed removal also can increase air movement in garden plots that will decrease conditions that are favorable for disease development.

Always disinfest machinery and other tools with steam, hot water under pressure, or a 10-percent solution of household bleach and water so pathogens do not spread from one area to another.

Crop rotation

Avoid planting the same crop in the same area of the garden year after year. Continuous culture of the same kind of crop provides an opportunity for pathogen buildup.

For example, rotate leafy vegetables with grains or corn, or rotate annuals or biennials in seed and flower beds. It is best to grow the same or closely related plants in the same soil only once in every three to five years. This practice "starves out" most pathogens that cause leaf, flower and stem diseases.

Crop rotation is not as effective against soilborne organisms, those pathogenic fungi, bacteria and nematodes that persist in the soil for up to 10 years or more.

Soil sanitation treatments

Occasionally disease-causing organisms that live in the soil may build up and prevent satisfactory growth of plants. Pathogen-free soil is desirable for houseplants, transplants and garden plots. Sterile potting mixes are available at many garden centers; however, it may be desirable to sanitize small quantities of soil on your own.

There are several nonchemical methods available to eradicate or reduce pathogens in the soil. The use of dry, steam or solar heat are the most effective nonchemical means to sanitize soil. The time to treat soil is before seeding or transplanting. Soil to be treated must be easily crumbled and without clods or large pieces of plant debris. Any amendments (manure, compost, peat moss, etc.) **must be added before treatment.** Soil also must have proper moisture. To test for this, gently squeeze a handful of soil. When the hand is opened, the soil ball should break apart somewhat. If it doesn't and the ball cannot be broken apart by gently pushing down on top of the ball, the soil is too wet.

Oven Sterilization. Place soil evenly but not more than 4 inches deep in non-plastic containers, such as seed flats, clay pots and glass or metal baking

Remove plant debris or infected plant parts after each growing season.

Eradication of weeds can break the life cycle of a pathogen and control it.

Sterile potting mixes are available at many garden centers; however, it may be desirable to sanitize small quantities of soil on your own.

Effective soil sanitation treatments include:

- **Oven sterilization**
- **Pressure-cooker sterilization**
- **Steam sterilization without pressure**
- **Soil solarization**

Create an environment unfavorable to pathogens.

pans. Tightly cover each container with aluminum foil. Insert a meat or candy thermometer through the foil into the center of the soil. Set the oven temperature between 180 and 200 F.

Heat the soil to at least 180 F and allow it to remain at this temperature for 30 minutes. Do not allow the temperature to go above 200 F since this may cause products toxic to plants. After heating, cool and remove containers from the oven. Leave the aluminum foil in place until ready for use. The heated soil will give off an odor. Microwaves or outdoor cookers also can be used.

Pressure-cooker sterilization. Pour several cups of water into the cooker. Place no more than 4 inches of soil in shallow containers on a rack out of the water. Level the soil, but do not pack it down. Cover each container with aluminum foil. Stack the containers to allow steam circulation. Close the lid, but leave the steam valve open somewhat until all the air is forced out and steam begins to escape. Then close the steam valve and heat at 10 pounds pressure for 15 minutes. Turn off the heat, allow the containers to cool, and remove. Again, leave the aluminum foil in place until you are ready to use the soil.

Steam sterilization without pressure. Pour about one inch of water into the sterilizing container. Follow the soil-preparation procedures listed earlier. Place filled soil containers on a rack that will hold them out of the water. Close the lid and bring the water to a boil. Open the lid just enough to prevent pressure from building up. When the steam begins to escape, continue boiling for 30 minutes. Turn off the heat and replace the lid. Remove the soil when cool.

Soil solarization. This method uses the sun's energy to heat small areas of soil to temperatures that are lethal to many soilborne organisms and weed seeds. Treat during the summer when there are high air temperatures and intense solar radiation. Loosen soil in the area that will be treated with a Rototiller or a shovel, wet and cover with a thin, clear polyethylene (plastic) film. Seal the edges of the plastic sheet with soil to prevent heat loss and retain moisture. Leave the plastic in place for several weeks. The longer the soil is exposed to the heat generated by the solarization process, the greater the kill of undesirable organisms. For more information, contact the Colorado State University Cooperative Extension Office in your county.

Avoiding toxicity from heated soil. With heavier soils and soils that contain a large amount of organic matter (manure, compost or peat moss), a toxic effect from heat sanitation may occur. This can cause poor seed germination, plant growth abnormalities or plant death. The toxicity is caused by an accumulation of ammonium compounds, soluble organic compounds, minerals or salts when the soil is heated too long or at too high a temperature. If soil toxicity is a problem, heavy irrigation of the treated soil may leach out many of these substances. Storing the soil two to three weeks without a cover will also reduce soil toxicity.

Cultural Management

Cultural management involves avoiding the onset of disease. To accomplish this without the use of pesticides, create an environment unfavorable to pathogens. For example:

1. Don't work in the garden when plants and soil are wet. Spores and cells of disease-causing organisms can spread from one plant to another and initiate new disease. Wet soils are easily compacted, which can decrease the amount of oxygen in the soil available for root uptake.
2. Make sure plants are spaced properly. Air movement is decreased when plants are grown too close together, and this allows moisture to remain on leaves for longer periods of time. Wider spacing in beds and landscape plantings promotes rapid drying after wet periods and stops development of foliage, flower and fruit pathogens.

3. Avoid excessive soil moisture. Over-watering enhances seed decay, damping-off and root rot diseases. Try not to plant in areas that have poor drainage or where water stands for several days following rains.

4. Fertilize plants properly based on soil nutrient analyses, using either organic or commercially prepared (inorganic) fertilizers.

In other words, use good cultural management. Healthy plants are less likely to have disease problems than weak, undernourished ones. Grow plants under optimum conditions and there will be fewer disease problems.

Control of most plant diseases **can** be accomplished without the use of pesticides. As long as sound cultural practices, sanitation and well-adapted plant varieties are used, there will be few disease problems. It is important to realize that you must accept some disease loss. Don't expect a "perfect" garden or plant if you do not want to use chemicals.

RECEIVED

JUL 08 1938

STATE PUBLICATIONS
Colorado State Library