


NR6.2/B53/1988/2

c.2




COLORADO BIRD DISTRIBUTION LATILONG STUDY



PHOTO BY LEONARD LEE RUE III

MAP OF COLORADO



COLORADO BIRD DISTRIBUTION
LATILONG STUDY

BY
THE COLORADO DIVISION OF WILDLIFE
(In Cooperation With The Colorado Field Ornithologists)

Editor:

Hugh E. Kingery

Assistant Editor:

Mary Beth Dillon

Records as of December 31, 1987

Editors, 1982 edition:

Steven J. Bissell

Charles Chase, III

Walter D. Graul

Hugh E. Kingery

Editors, 1978 edition:

Hugh E. Kingery

Walter D. Graul

Assistant Editors, 1978 edition:

David Alles Barry Knapp

Bob Andrews Mildred Snyder

Copyright, Colorado Division of Wildlife, 1988

Send inquiries about obtaining copies to:

Colorado Division of Wildlife
6060 Broadway
Denver, CO 80216

INTRODUCTION

This report synthesizes the contributions of over 300 field ornithologists, and summarizes the present knowledge of distribution, abundance, and habitat preferences of the birds of Colorado. It presents the information on a regional basis, using a plotting system based on latitudes and longitudes--"latilong" blocks. Using this guide, the reader may determine the status of a particular species in a given latilong block, as well as its relative abundance and the habitats in which it can be expected to occur.

The latilong plotting system follows the method used by Skaar¹ to report Montana bird distribution. Latitudes and longitudes divide Colorado into 28 blocks roughly 50 X 70 miles. (See the master map on the inside of the front cover; for more accuracy, consult an official Colorado highway map issued by the State Highway Department). Thus, on the attached species sheets, the dashes correspond to the latilong blocks and represent a map of Colorado.

For three basic reasons, this system differs from traditional ways of presenting distributional information. First, the system allows for the presentation of much detailed information in a small amount of space. Second, it allows for conformity on a national and international basis. For instance, the U.S. breeding bird roadside census, the international breeding bird plot census and all banding data are based on a latitude-longitude coordinate system. Third, the coordinate system lends itself to computerization; it permits efficient storage, retrieval, and amendment of the information.

The arbitrary nature of the latilong boundaries, like state boundaries, does create some unexpected effects. For example, latilong 4 includes Boulder, Longmont, and Fort Collins, a myriad of reservoirs and farm ponds, much irrigated farmland, short-grass prairie, the Rawhide power plant reservoir, and mountain habitats from pinon pine near Fort Collins to alpine tundra along the Continental Divide. This block ranges from 4,860 feet on the Poudre River to 14,256 feet at the summit of Longs Peak. Thus to plot the actual distribution of species within a single latilong block, the user must combine habitat data cited here with actual habitats within the block.

For this reason, the Breeding Bird Atlas, now underway, promises to refine our knowledge of the breeding birds of Colorado. Like the Latilong system, the Atlas will present information in conformity with a nationally-used scheme--the scale and data codes used by atlases in other states. Furthermore, the data will fit into the existing latilong scheme. At this time Atlas data is preliminary, so that this edition includes only a handful of records from Atlas field work.

¹Skaar, P.D. 1975. Montana Bird Distribution. Published by the author, Bozeman, MT.

RESULTS

We list herein 427 species that have been recorded in Colorado within the past 22 years (since the 1965 publication of Bailey and Niedrach's seminal work, "Birds of Colorado"). This edition includes records through December 31, 1987, plus three 1988 new breeding species in Colorado (Little Blue Heron, Least and Great Crested Flycatcher) and one new species for the state (Ruffed Grouse) documented by a 1988 specimen. The following 15 species, (which complete recorded state list of 442 species), were recorded before 1965 and are not included in the species maps which follow..

| | | |
|-------------------|--------------------|-------------------------|
| Brown Pelican | Eskimo Curlew | Alder Flycatcher |
| Anhinga | Long-tailed Jaeger | Dusky-capped Flycatcher |
| Reddish Egret | Ivory Gull | Long-billed Thrasher |
| Roseate Spoonbill | Ancient Murrelet | Lucy's Warbler |
| Wood Stork | Barred Owl | Cerulean Warbler |

Observers who find these species, or any others not listed in the main section, should document them with copious detail on the form provided in Appendix III.

We stress that the information in the species listings includes only data actually submitted--we made no assumptions. (For species marked with an asterisk, only records approved by the Colorado Field Ornithologists Official Records Committee (CFO) are included. This also applies to certain species for which CFO has only recently dropped its documentation requirement, and to some other species out of their normal range or difficulty to identify. At the time of this edition, CFO had reviewed records submitted to it through December 31, 1986.) Using only actual data has resulted in some inconsistent distributional patterns. For instance, we may have documentation that a particular species is a year-round resident in several latilong blocks, but in adjacent blocks have documentation of breeding only. Some such differences may simply represent gaps in our knowledge, although others may be real and based on habitat difference. Therefore use data for a given species cautiously, and employ all data including, especially, habitat limitations.

Each edition of this publication serves as a springboard for gathering more complete and refined information in the future. This continuing project relies directly on the additions and corrections we continue to receive. Consequently we encourage all persons doing field work on Colorado birds to submit new information. Appendix I provides a form for reporting new information on latilong status, abundance, and habitat preference for one species. Use the Appendix II form if you have several changes which do not require extra documentation. For rare species (those with an asterisk, those not listed, those out of season, or those out of range) use Appendix III.

The third edition of the "Latilong Study" has incorporated 1,760 changes in status and habitat. We stress that we need much more documentation of habitat preferences, as well as better distributional information. Data on common or uncommon species can be as valuable, or even more valuable, as that on rare species. Please send data which you gather, using the appropriate form(s), to: Colorado Bird Atlas, c/o Zoological Collections, Denver Museum of Natural History, City Park, Denver, CO 80205.

ACKNOWLEDGMENTS

We express special thanks to all those who contributed time, thought, and effort in submitting records and helping to put together this document. Contributors have used special diligence and care in reporting data. The first edition listed all the contributors, but the second did not, partly because of the large number and partly because we might accidentally omit someone. Essentially, almost every active bird watcher in Colorado has contributed to this study. Contributors now number at least 300, and the Editors of all three editions acknowledge each contributor with appreciation.

We add our special thanks to Patty Frede, for her extra attention and patience with the editor and for her high caliber word processing expertise.

The Colorado Division of Wildlife has supported this edition with a grant, manuscript preparation, and printing. Copies of the worksheets which form the data background for this edition are on file with the Division, the Denver Museum of Natural History (Library), and the Colorado Bird Atlas Partnership.

REQUEST FOR DATA

In keeping with the cooperative nature of this publication, the Division and CFO issue this Request for Data.

Please report observations of the following:

- If you obtain photographs or a specimen.

* - Any observation -- use the Appendix III form.

Acc - If you observe in any designated "Accidental" latilong. If the species has no # or *, use the Appendix I or II form.

Blank -If a latilong is blank, and the species has no * or #, use the Appendix I or II form.

Please send to Colorado Bird Atlas, c/o Zoology Department, Denver Museum of Natural History, 2001 Colorado Blvd., Denver, CO 80205.

THE FORMAT

The format uses several codes in order to report the information. There follows an explanation of the Status codes, the Habitat listings, and the Abundance factors. Additionally, for your convenience the status and abundance symbols are given on each species page.

STATUS

In each latilong block the symbols listed below are used to indicate a species' status. The categories override each other in the order given. For example, if a species both breeds in and migrates through a given latilong block, it would be listed as B for that block.

R - Resident (Breeds and present year-round).

N - Non-breeder present year-round, or a year-round resident whose breeding has not been documented.

B - Breeding (Definite evidence of nesting, i.e. nests or flightless young observed).

b - Very likely breeding (Based on the observer's opinion. For example, birds observed regularly in suitable nesting habitat during the nesting season).

n - Non-breeder, but present during the nesting season.

W - Winter visitor (Present during most or all of the period from December 16-February 28).

M - Migrant (Definition from O.W. Pettingill¹ - any movement between two areas).

(In any latilong which has less than four records - listed marginally as "Acc" - i.e. Accidental)

An example may help clarify the system. If we have reports of Mountain Bluebirds nesting in South Park (Block 11); migrating in Greeley, Denver, the Black Forest, and Pueblo (Blocks, 5, 12, 19); and wintering around Cortez, Durango, and Trinidad (Blocks 22, 23, 26), and two records from Bonny dam, the map would look like this:

| | | |
|------------|--|---|
| Acc. in 14 | <u> </u> <u> </u> <u> </u> <u>B</u> <u>M</u> <u>M</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>M</u> <u> </u> <u>W</u> <u>W</u> <u> </u> <u> </u> <u>W</u> <u> </u> | (We left a space down the middle of the state to help orient you to the map.) |
|------------|--|---|

¹Pettingill, Olin Sewall, Jr. 1970. Ornithology in Laboratory and Field, Fourth edition. Burgess Publishing Co., Minneapolis, MN. p. 267.

KEY TO SYMBOLS

- # - Species recorded in the state only on the basis of sight records. C.F.O. Records Committee solicits photographs and substantiating data.
- * - Species for which Colorado has no more than ten records. Please document using the form in Appendix III.
- Acc - "Accidental" in the specified Latilong(s). These are species with fewer than four records in a Latilong.

ABUNDANCE

This aspect of biology, a somewhat subjective problem, we originally intended to disregard. To a degree, the extent of distribution will indicate the relative abundance of a species. However, whereas the Horned Lark and Mountain Plover show similar distributions on the eastern plains, their relative abundance differs markedly. We used the following terms in attempting to record abundance values.

- Ab - Abundant: Breeding - at least 1 per acre, e.g. Horned Lark, Lark Bunting, Brewer's Sparrows, and in many areas Red-winged Blackbirds. Migration and Winter - counts of 25 per day usual.
- C - Common: Breeding - at least 1 per 5 acres, e.g. Meadow-lark, Western Grebes in marshes. Migration and Winter - counts of 10 per day usual.
- FC - Fairly Common: Breeding - less than 1 per 5 acres, but can usually find in appropriate habitat in any one day. Migration and Wintering - one or more per field trip.
- U - Unusual: See no more than 5 during the season in a given latilong.
- R - Rare: Fewer than 10 records in a latilong.
- Und - Undetermined: Not enough information to classify.
- Irr - Irregular: Abundant at times; extremely scarce at other times.
- Acc - Accidental or straggler (1-3 records in a latilong block). If a species has this designation, it may just indicate that we have too few data to classify it under one of the other status categories.

HABITAT

Clarifying Description

| Code | Name | Dominant Vegetation Type | Some Associated Plants |
|-------------------------------|-------------------|-----------------------------------|--|
| <u>A - Alpine</u> | | | |
| T | Tundra | | |
| AT | Alpine Transition | KRUMMHOLZ EVERGREEN | |
| <u>CF - Coniferous Forest</u> | | | |
| Lm | Limber Pine | LIMBER PINE | whitebark pine, whiproot clover, common juniper |
| SF | Spruce-Fir | ENGELMANN SPRUCE SUBALPINE FIR | heartleaf arnica, twinflower, grouse whortleberry, buffaloberry |
| LP | Lodgepole Pine | LODGEPOLE PINE | kinnikinnick, common juniper, buffaloberry, Rocky Mtn. & grouse whortleberry |
| Bc | Bristlecone Pine | BRISTLECONE PINE | whiproot clover, blue-joint reedgrass |
| PP | Ponderosa Pine | PONDEROSA PINE | Rocky Mtn. & common juniper, antelope bitterbrush, mtn. mahogany, Arizona, Idaho & spike fescue, common snowberry, side-oats grama |
| DF | Douglas-fir | DOUGLAS-FIR | mtn. mahogany, mtn. & common snowberry, cliff jonesia, Idaho fescue |
| PJ | Pinyon-Juniper | PINYON PINE UTAH JUNIPER | blue grama, big sagebrush, gambel oak, mtn. mahogany, antelope bitterbrush, western & bluebunch wheatgrass |

Clarifying Description

| Code | Name | Dominant Vegetation Type | Some Associated Plants |
|------------------------|--|----------------------------------|---|
| <u>GL - Grasslands</u> | | | |
| SgP | Shortgrass-Prairie | BLUE GRAMA-BUFFALOGRASS | western wheatgrass, sun sedge (as on Pawnee Nat'l Grassland) |
| CG | Cholla Cactus | CHOLLA CACTUS | galleta, red threeawn, ring muhly, needle and thread grass, yucca (as in northcentral and southcentral Pueblo County) |
| | Grassland | BLUE GRAMA | |
| SgSD | Shortgrass-Semi- Desert | GRAMA-GALLETA | sand bluestem, little bluestem, big sagebrush, loco, fourwing saltbush, side-oats grama, hairy grama |
| MXP | Mixed grasses of habitat alteration areas on Plains (plantings & intro) | CRESTED WHEATGRASS THISTLE | side-oats grama, little bluestem |
| TgP | Tallgrass Plains | LITTLE BLUESTEM-SANDSAGE | switchgrass (as on Yuma Sandhills) |
| SgM | Shortgrass-Mountains | ARIZONA FESCUE MOUNTAIN MUHLY | rubber rabbitbrush, horsebrush (as in San Luis Valley and South Park) |
| SS | Sand Sage Prairie | SAND SAGEBRUSH-YUCCA | blue grama, buffalo grass, side-oats grama, red threeawn (as on Comanche Nat'l Grassland in Baca County) |
| MmP | Mountain Meadow- Parkland (wet & dry) | TUFTED HAIRGRASS | alpine timothy, Parry oatgrass, sedge sp. |

Clarifying Description

| Code | Name | Dominant Vegetation Type | Some Associated Plants |
|-----------------------------|--|--|---|
| <u>S - Shrublands</u> | | | |
| SaRb | Sagebrush-Rabbitbrush (of shrub steppe) | BIG SAGEBRUSH RABBITBRUSH | |
| DS | Greasewood-Sagebrush or Greasewood-Four-wing Saltbush | GREASEWOOD-BIG SAGEBRUSH GREASEWOOD-FOURWING SALTBUSSH | (as in San Luis Valley) |
| Sa | Sagebrush (of Mtn. shrubs) | BIG SAGEBRUSH | antelope bitterbrush, needle and thread grass, Idaho & Thurber fescue, Indian ricegrass |
| Sc | Scrub Oak (of Mtn. shrubs) | GAMBEL OAK | mtn. mahogany, snowberry, Utah and Saskatoon serviceberry |
| MM | Mountain Mahogany (of Mtn. shrubs) | MOUNTAIN MAHOGANY | bluebunch wheatgrass, needle and thread grass, skunkbush, Idaho fescue |
| <u>R - Deciduous Forest</u> | | | |
| As | Aspen | ASPEN | mtn. snowberry, silvery lupine, Kentucky bluegrass, Thurber fescue |
| RpL | Riparian Lowland (including edges of lakes) | PLAINS COTTONWOOD WILLOW-TAMARISK | snowberry, buffaloberry, wild-rye, sedges (below 6,000 ft.) |
| RpT | Riparian Transition | NARROWLEAF COTTONWOOD WILLOW-ALDER | Douglas-fir, timothy, sedge sp., blue spruce, (6,000 to 9,000 ft.) |
| RpH | Riparian Highland | WILLOW-ALDER | marsh marigold, tufted hairgrass, cinquefoil (above 9,000 ft.) |

Clarifying Description

| Code | Name | Dominant Vegetation Type | Some Associated Plants |
|---------------------|--------------------------------|---|--|
| <u>AQ - Aquatic</u> | | | |
| Ms | Marsches, bogs, wet hummocks | CATTAIL-BULRUSH | may be found in conjunction with lake or reservoir (specify) |
| W/OG | Wet open ground | ALKALI FLATS, MUD FLATS, GRAVEL BEDS | (specify) |
| Ri | Open Water-Streams and Rivers | | |
| L | Open Water-Lakes or Reservoirs | | |
| Miscellaneous | | | |
| Ag | Agricultural Areas | ORCHARDS, SHELTER BELTS, DWELLINGS, TREE FARMS | (specify) |
| Cr | Croplands | ANNUAL CROPS | wheat, corn, milo, etc. |
| U | Urban | CITIES, TOWNS (including parks within cities) | |
| O | Other | BRIDGES, CAVES, ATTICS, CLIFFS, BANKS, SAND DUNES, TALUS SLOPES | (specify) |

A LATILONG GAZETTEER

We have noted that records from some locations which lie close to a latilong boundary may have been reported from an incorrect Latilong. Part of this stems from gathering information for the 1978 edition, in which the latitudes and longitudes were not shown precisely. In any event, the list below identifies latilongs for a few of these sometimes misplaced locations.

| | | | |
|--------------------------------|-------|----------------------|----|
| Antero Res. | 18 | Harts Basin Res. | 15 |
| Baseline Res. (mostly) | 4 | Kinney Lake | 20 |
| Boulder (split by Baseline Rd) | 4/11 | Russell Lakes S.W.A. | 24 |
| Brush Hollow Res. | 18 | Rye | 26 |
| Denver (split by I-25) | 11/12 | Sweitzer Lake | 16 |
| Flagler State Wildlife Area | 14 | | |

| | | | | | |
|-------------------------------------|----------------------------------|------------|-----------------|----------|---------|
| <u>Red-necked Grebe*</u> | <u>Podiceps grisegena</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 18, 19, 21, 25 | | | M M | L | |
| | | | — — M M — — | | |
| | | | — — M M — — | | |
| | | | — — M — — | | |
| | | Abundance: | | R | |
| <u>Eared Grebe</u> | <u>Podiceps nigricollis</u> | Habitat: | Breeding | Migr. | Winter |
| | | | B M B b B M M | Ms,L | Ms,L |
| | | | M b B B b b M | | |
| | | | B n B B b b b | | |
| | | | B M B B B — M | | |
| | | Abundance: | FC | C-Ab | |
| <u>Western Grebe</u> | <u>Aechmophorus occidentalis</u> | Habitat: | Breeding | Migr. | Winter |
| | | | n M B R B n M | Ms,L | L,Ri,Ms |
| | | | n M M n B n M | | |
| | | | M B M B B B n | | |
| | | | n n b B B M M | | |
| | | Abundance: | FC | FC-Ab | R |
| <u>Clark's Grebe</u> | <u>Aechmophorus clarkii</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 7, 9, 10, 12, 18, 19, 21 | | | M M M M b B — M | L, Ms | L |
| | | | M B — M B B M | | |
| | | | — B — — — — | | |
| | | Abundance: | U | U | |
| <u>American White Pelican</u> | <u>Pelecanus erythrorhynchos</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 2, 8, 9, 22, 26 | | | M M M n B M M | L | Ms,L |
| | | | M M n M n — M | | |
| | | | M M M n M n M | | |
| | | | M M n M M M M | | |
| | | Abundance: | (one place) | R-Ab | |
| <u>Double-crested Cormorant</u> | <u>Phalacrocorax auritus</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 2, 9, 10, 16, 22, 23 | | | M M M B B B B | L,Ri,RpL | L |
| | | | M M M B B B b | | |
| | | | M M M B b B n | | |
| | | | M M M M n M M | | |
| | | Abundance: | FC | FC-Ab | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|---|--|------------------------|------------------------|---------|
| <u>Snowy Egret</u> | | Breeding | Migr. | Winter |
| <u>Egretta thula</u> | M n B b B b M | Habitat: Ms, RpL | Aq | |
| Acc. in 9 | <u>B</u> <u>M</u> <u>M</u> <u>b</u> <u>B</u> <u>b</u> <u>M</u> | | | |
| | <u>b</u> <u>b</u> <u>b</u> <u>M</u> <u>M</u> <u>B</u> <u>M</u> | | | |
| | <u>M</u> <u>M</u> <u>B</u> <u>B</u> <u>B</u> <u>M</u> <u>M</u> | | | |
| | | Abundance: U-FC | FC | |
| <u>Little Blue Heron</u> | | Breeding | Migr. | Winter |
| <u>Egretta caerulea</u> | M M M M M M - | Habitat: Aq, RpL | | |
| Acc. in 1, 6, 8, 10, 11, 15, 18, 21, 25, 28 | <u>M</u> <u>-</u> <u>M</u> <u>M</u> <u>M</u> <u>-</u> | | | |
| | <u>M</u> <u>-</u> <u>-</u> <u>M</u> <u>M</u> <u>M</u> | | | |
| | <u>-</u> <u>-</u> <u>B</u> <u>M</u> <u>-</u> <u>-</u> <u>M</u> | | | |
| | | Abundance: R | R | |
| <u>Tricolored Heron</u> | | Breeding | Migr. | Winter |
| <u>Egretta tricolor</u> | M M M M M M - | Habitat: Aq, RpL | | |
| Acc. in 4, 5, 12, 19, 20 | <u>-</u> <u>-</u> <u>-</u> <u>M</u> <u>M</u> <u>-</u> | | | |
| | <u>-</u> <u>-</u> <u>-</u> <u>M</u> <u>M</u> <u>-</u> | | | |
| | <u>-</u> <u>-</u> <u>-</u> <u>-</u> <u>-</u> <u>-</u> | | | |
| | | Abundance: R | R | |
| <u>Cattle Egret</u> | | Breeding | Migr. | Winter |
| <u>Bubulcus ibis</u> | M M M n B M M | Habitat: Ms, RpL | Ms, Ri, L, Cr | |
| Acc. in 1, 2, 6, 9, 10, 21, 22, 23, 27, 28 | <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>B</u> <u>M</u> <u>M</u> | | | |
| | <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | |
| | <u>M</u> <u>M</u> <u>B</u> <u>-</u> <u>M</u> <u>M</u> <u>M</u> | | | |
| | | Abundance: U | U | |
| <u>Green-backed Heron</u> | | Breeding | Migr. | Winter |
| <u>Butorides striatus</u> | M M M n M M B | Habitat: RpL, Ms | Ms, RpL, RpT, L, Ri | RpL, Ri |
| Acc. in 1, 2, 3, 8, 15, 16, 22, 23, 26 23: one wintered, 1976 | <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>b</u> <u>B</u> <u>b</u> | | | |
| | <u>M</u> <u>W</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | |
| | | Abundance: R, U-plains | R | |
| <u>Black-crowned Night-Heron</u> | | Breeding | Migr. | Winter |
| <u>Nycticorax nycticorax</u> | B B B B B b B | Habitat: Aq, RpL | Aq, RpL | |
| | <u>B</u> <u>B</u> <u>B</u> <u>R</u> <u>B</u> <u>-</u> <u>M</u> | | | |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> | | | |
| | <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>M</u> | | | |
| | | Abundance: FC | FC | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|---------------------------------------|--|----------------------------------|------------------------|---------------|
| <u>Yellow-crowned Night-Heron</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Nycticorax violaceus</u> | M M B M M M M | Aq,RpL | Aq,RpL | |
| Acc. in 4, 11, 19, 20, 21, 28 | | | | |
| | | Abundance:R | R | |
| <u>White Ibis*</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Eudocimus albus</u> | | Habitat: | | |
| Acc. in 21 | M | | | |
| | | Abundance:R | R | |
| <u>Glossy Ibis*#</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| | Habitat: | | | |
| Acc. In 16 | M | | | |
| | | Abundance:R | R | |
| <u>White-faced Ibis</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Plegadis chihi</u> | M M M b M M M M M M b M M M B B | B M M M M M M M M M M M | Ms | Aq,Ag |
| Acc. in 13 | | Habitat: | | |
| | | Abundance:U | U-C | |
| <u>Black-bellied Whistling-Duck*#</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Dendrocygna autumnalis</u> | M | Habitat: | L, 1 record only | |
| Acc. in 11 | | | | |
| Record is possible escape | | | | |
| | | Abundance:Und | Und | |
| <u>Tundra Swan</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Cygnus columbianus</u> | M M M M M M M M M M M M | M M M M M M M M | L,Ms | |
| Acc. in 9, 15, 17, 18, 21, 22, 28 | | Habitat: | | |
| | | Abundance:U | U | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|--|--|----------------|--|-----------------|
| <u>Trumpeter Swan*</u> <u>Cygnus buccinator</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 11, 12, 18, 20 | M — — M M — — — — — M — M — — — — — — — | | L | |
| | | Abundance: | R | |
| <u>Greater White-fronted Goose</u> <u>Anser albifrons</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 9, 15, 16, 21, 25 | W M — W M M M W M — — M M M — M M M — M M M — M M M — M M M | | L,Ms,Cr | L,Cr |
| | | Abundance: | R-U | R |
| <u>Snow Goose</u> <u>Chen caerulescens</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 16, 17 | M M M W M M M W M M W W — M — M M M M W W — M M M M W W | | L,Ms,Cr | L,Cr |
| | | Abundance: | Ab-SE Colo U-SE Colo U-Rest of R-Rest of State State | |
| <u>Ross' Goose</u> <u>Chen rossii</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 6, 14, 15, 16, 17, 20, 23, 25, 28 | W — — M M M M M M — M M M — M — M — M | | L,RI,RpL, Cr | L |
| | | Abundance: | R | R |
| <u>Brant*</u> <u>Branta bernicla</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 4, 12 | — — — M M — — — — — — — — | | L | |
| | | Abundance: | R | |
| <u>Canada Goose</u> <u>Branta canadensis</u> | Habitat: | Breeding | Migr. | Winter |
| | R R B R R R R R R R R R W R R R B B W W W R M R R W N W | Aq,RpL, MXP | Aq,Cr, RpL,MXP | Aq,Cr,MXP, U |
| | | Abundance: | C-Ab | C-Ab Ab |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|--|--|----------------------------------|-------------------|--------|
| Wood Duck | | Breeding | Migr. | Winter |
| <u>Aix sponsa</u> | M b M R R M B R M M M M M M M M | B B b R — B B M M M — M | Habitat: Ri,L,RpL | Aq,RpL |
| Acc. in 1, 3, 9, 15, 16, 17, 18, 21, 25, 28 | | | L | |
| | | Abundance: U-R | U-R | R |
| Green-winged Teal | | Breeding | Migr. | Winter |
| <u>Anas crecca</u> | R R B R R R B N B R R B b R R R | R R R R — R R M W b M W | Habitat: Aq | Aq |
| | | | Aq | Aq |
| | | Abundance: FC | Ab | FC |
| American Black Duck | | Breeding | Migr. | Winter |
| <u>Anas rubripes</u> | — M M M — M M — M — M — M — — — — | Habitat: L | | |
| Acc. in 3, 4, 5, 11, 12, 14, 19 | | | | |
| | | Abundance: R | | |
| Mallard | | Breeding | Migr. | Winter |
| <u>Anas platyrhynchos</u> | R R R R R R R R R R R R R R R R | R R R R B R R R R R R R | Habitat: Aq,Cr,R | Aq,Cr |
| | | | Aq,Cr | Aq,Cr |
| | | Abundance: C | Ab | C-Ab |
| Northern Pintail | | Breeding | Migr. | Winter |
| <u>Anas acuta</u> | R b B R W B M R W M R b W B R R | R R R R B b R R b b M W | Habitat: Aq | Aq |
| | | | Aq | Aq |
| | | Abundance: FC | Ab | C |
| Blue-winged Teal | | Breeding | Migr. | Winter |
| <u>Anas discors</u> | B B B B M B B B n b b b M M B B | B B B B M B B B b B b B | Habitat: Aq | Aq |
| | | | Aq | Aq |
| | | Abundance: FC | Ab | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|--------------------------------------|----------------|----------------|---------------|-------------------|----------------|-----------|
| <u>Cinnamon Teal</u> | B B B B | b B b | Habitat: | Breeding Aq, W/OG | Migr. Aq, W/OG | Winter |
| <u>Anas cyanoptera</u> | <u>B B B B</u> | <u>B M b</u> | | | | |
| Acc. in 13, 21 | <u>B B b b</u> | <u>B b M</u> | | | | |
| | <u>b B B B</u> | <u>b B M</u> | | | | |
| | | | Abundance: FC | R-C | | |
| <u>Northern Shoveler</u> | R M B R | b b b | Habitat: | Breeding Aq | Migr. Aq | Winter L |
| <u>Anas clypeata</u> | <u>N M M R</u> | <u>R M W</u> | | | | |
| | <u>M n b b</u> | <u>b R M</u> | | | | |
| | <u>M M B B</u> | <u>B M b</u> | | | | |
| | | | Abundance: U | Ab | U | |
| <u>Gadwall</u> | R b B R | B R R | Habitat: | Breeding Aq | Migr. Aq | Winter Aq |
| <u>Anas strepera</u> | <u>M b M R</u> | <u>R b b</u> | | | | |
| | <u>M B R b</u> | <u>R R b</u> | | | | |
| | <u>M B R R</u> | <u>b M M</u> | | | | |
| | | | Abundance: FC | Ab | FC | |
| <u>Eurasian Wigeon#</u> | | | Habitat: | Breeding L | Migr. | Winter |
| <u>Anas penelope</u> | <u>— — M M</u> | <u>— — M M</u> | | | | |
| Acc. in 4, 11, 12, 15, 19, 24, 26 | <u>— — M —</u> | <u>— — M —</u> | | | | |
| | <u>— — M —</u> | <u>— — M —</u> | | | | |
| | | | Abundance: R | | | |
| <u>American Wigeon</u> | R B B R | R b W | Habitat: | Breeding Aq | Migr. Aq | Winter L |
| <u>Anas americana</u> | <u>W b R R</u> | <u>R M M</u> | | | | |
| | <u>M n b M</u> | <u>R R M</u> | | | | |
| | <u>M B R B</u> | <u>b M b</u> | | | | |
| | | | Abundance: C | Ab | FC | |
| <u>Canvasback</u> | B M B b | b M M | Habitat: | Breeding Ms, L | Migr. L | Winter L |
| <u>Aythya valisineria</u> | <u>W M M W</u> | <u>N M M</u> | | | | |
| Acc. in 10 | <u>W M M M</u> | <u>M M n</u> | | | | |
| | <u>M M B B</u> | <u>B M M</u> | | | | |
| | | | Abundance: R | C | R | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | |
|--|--|------------|---|
| <u>Redhead</u> <u>Aythya americana</u> | R b B R B b b W M M R R M M M M B M B R b B B B B B M M | Habitat: | Breeding L,Ms Migr. L,Ms Winter L |
| | | Abundance: | FC-C Ab U-FC |
| <u>Ring-necked Duck</u> <u>Aythya collaris</u> | R b B R M M M N R B R W — M R b B M M M M M R M M M M M | Habitat: | Breeding L,Ms,RpT Migr. L,Ms Winter L |
| Acc. in 5, 6, 8, 23, 24, 25, 26 | | Abundance: | R FC-C U |
| <u>Greater Scaup</u> <u>Aythya marila</u> | M — M M M M M — — — M — M — M M M M — — | Habitat: | Breeding L Migr. L Winter L |
| Acc. in 12 | | Abundance: | R |
| <u>Lesser Scaup</u> <u>Aythya affinis</u> | b M B R M M M W R M W W M M W M M M M W M M M B b b M M | Habitat: | Breeding L Migr. L Winter L |
| Acc. in 12 | | Abundance: | R Ab U |
| <u>Harlequin Duck*</u> <u>Histrionicus histrionicus</u> | — — — — M — — — — — — — — — — — — — — — — | Habitat: | Breeding L Migr. L Winter L |
| Acc. in 12 | | Abundance: | R |
| <u>Oldsquaw</u> <u>Clangula hyemalis</u> | M M M M M M M M M — M M — M M — M M M M M — M — — M — — | Habitat: | Breeding L,Ri Migr. L Winter L |
| Acc. in 1, 3, 6, 7, 8, 9, 14, 15, 18, 20, 21, 23, 26 | | Abundance: | R |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|--------------------------------|-----------------|----------|-------|-----------|
| Black Scoter | | Breeding | Migr. | Winter |
| <u>Melanitta nigra</u> | | | L | |
| Acc. in 6, 8, 16 | | | | |
| Fall records only | | | | |
| | Habitat: | | | |
| | M — — M M — | | | |
| | — M — — — — | | | |
| | — — — — — — | | | |
| | Abundance: | R | | |
| Surf Scoter | | Breeding | Migr. | Winter |
| <u>Melanitta perspicillata</u> | | | L | |
| Acc. in 6, 8, 16, 19, | | | | |
| 20, 23 | | | | |
| | Habitat: | | | |
| | M — — M M — | | | |
| | — M — — M M — | | | |
| | — M — — — — | | | |
| | Abundance: | R | | |
| White-winged Scoter | | Breeding | Migr. | Winter |
| <u>Melanitta fusca</u> | | | L | |
| Acc. in 1, 7, 8, 9, 15, | | | | |
| 16, 22, 23, 25, 26 | | | | |
| | Habitat: | | | |
| | M — — M M — M | | | |
| | — M — — — — M | | | |
| | — M — — — — — | | | |
| | Abundance: | R | | |
| Common Goldeneye | | Breeding | Migr. | Winter |
| <u>Bucephala clangula</u> | | | Aq | L,Ri |
| | Habitat: | | | |
| | W W W W W W M | | | |
| | W W W W W W — | | | |
| | W W M W W W W | | | |
| | — M W M W W W W | | | |
| | Abundance: | C C | | |
| Barrow's Goldeneye | | Breeding | Migr. | Winter |
| <u>Bucephala islandica</u> | | | L,CF | L,Ri L,Ri |
| Acc. in 1, 2, 6, 12, 16, 17, | | | | |
| 28 | | | | |
| | Habitat: | | | |
| | M M B W M — | | | |
| | W W W W W — — | | | |
| | — M M M M W — | | | |
| | — — — — W — M | | | |
| | Abundance: | R U R | | |
| Bufflehead | | Breeding | Migr. | Winter |
| <u>Bucephala albeola</u> | | | L,Ri | L,Ri |
| Acc. in 13 | | | | |
| | Habitat: | | | |
| | W M M W W W M | | | |
| | W M M W W M W | | | |
| | W M M M W M W | | | |
| | — M W M M M W W | | | |
| | Abundance: | C U | | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|-------------------------------|------------------------------|--|--------------------------|---------------|
| <u>Hooded Merganser</u> | <u>Lophodytes cucullatus</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| | | W B W W b | Habitat: Ms,RpL | L,Ri L,Ri |
| | | W M — W — | | |
| | | W M M M — | | |
| | | — W M W — | | |
| | | Abundance: R U R-U | | |
| <u>Common Merganser</u> | <u>Mergus merganser</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| | | B R R R W W W | Habitat: Aq,RpL | Aq Aq |
| | | R R R R — W W | | |
| | | R R R W W W W | | |
| | | b R R M b W W | | |
| | | Abundance: FC C C | | |
| <u>Red-breasted Merganser</u> | <u>Mergus serrator</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| Acc. in 9, 20 | | W M M M — | Habitat: Aq | Aq |
| | | — W — W — | | |
| | | M M — W — M | | |
| | | — — — W M — | | |
| | | M M M — M — | | |
| | | Abundance: C R | | |
| <u>Ruddy Duck</u> | <u>Oxyura jamaicensis</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| Acc. in 13 | | B M B B B b b | Habitat: Ms,L | Aq |
| | | M b M B B M B | | |
| | | M n B M B B b | | |
| | | B B B B b M M | | |
| | | Abundance: FC C-Ab | | |
| <u>Turkey Vulture</u> | <u>Cathartes aura</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| | | B B B B b M M | Habitat: Rpl,Rpt, cliffs | All |
| | | B B B B B M B | | |
| | | B b b b B B b | | |
| | | B B B B B B B | | |
| | | Abundance: FC | FC | |
| <u>Osprey</u> | <u>Pandion haliaetus</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| Acc. in 9 | | B B B B M M M | Habitat: R,L,Ri | L,Ri,RpL, RpT |
| | | M M b M M — b | | |
| | | b M b M M M M | | |
| | | M B M b M — M | | |
| | | Abundance: U | U | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|------------------------------------|---------|-------|------------------|-------------|------------|---------------------------|
| <u>Northern Goshawk</u> | R R R R | W W W | Habitat: | Breeding CF | Migr. CF,R | Winter CF,R,As, U,SaRb |
| <u>Accipiter gentilis</u> | RRB R | R B W | | | | |
| Acc. in 7, 21, 28 | R R B B | R M | | | | |
| | RRB R | R R M | | | | |
| | | | Abundance: U | U | U | |
| <u>Common Black-Hawk#*</u> | | | Habitat: | Breeding | Migr. | Winter |
| <u>Buteogallus anthracinus</u> | --- | M | | | RpL | |
| Acc. in 11 | --- | --- | | | | |
| | --- | --- | | | | |
| | | | Abundance: _____ | R | | |
| <u>Red-shouldered Hawk</u> | | | Habitat: | Breeding | Migr. | Winter |
| <u>Buteo lineatus</u> | M M M | M M M | | | RpL,U | |
| Acc. in 4, 5, 7, 11, 12, 14, 19 | --- | M | | | | |
| | --- | --- | | | | |
| | --- | --- | | | | |
| | | | Abundance: _____ | R | | |
| <u>Broad-winged Hawk</u> | | | Habitat: | Breeding | Migr. | Winter |
| <u>Buteo platypterus</u> | B M | M M M | | U, | RpL,Ag,U | |
| Acc. in 5, 13, 17, 20, 21, 27 | --- | M | 1 record | | | |
| | --- | --- | | | | |
| | --- | --- | | | | |
| | | | Abundance: R | R | | |
| <u>Swainson's Hawk</u> | | | Habitat: | Breeding | Migr. | Winter |
| <u>Buteo swainsoni</u> | B B B B | B B B | | GL,Ag,RpL, | R,S,A,GL, | |
| | BBB B | B B B | | RpT,DS | Ag | |
| | BBB B | B B B | | | | |
| | B M B B | B B B | | | | |
| | | | Abundance: FC | FC | | |
| <u>Red-tailed Hawk</u> | | | Habitat: | Breeding | Migr. | Winter |
| <u>Buteo jamaicensis</u> | R R R R | R R R | | CF,Ag,R,As | All | Ag,R,U |
| | RRR R | R R R | | | | |
| | RRR R | R R R | | | | |
| | RRR R | R R R | | | | |
| | | | Abundance: FC | FC | FC | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|--------------------------|---------|-------|------------|-----------------------|--------------------|-------------------------|
| <u>Ferruginous Hawk</u> | B B B N | R R B | Habitat: | Breeding SgP, Ag, MmP | Migr. GL, DS, Ag | Winter GL, Ag, RpL, RpT |
| <u>Buteo regalis</u> | R b M b | R R R | | | | |
| Acc. in 16, 22 | N M M B | R R R | | | | |
| | M W R B | B R R | | | | |
| | | | Abundance: | FC | U | U |
| <u>Rough-legged Hawk</u> | W W W W | W W W | Habitat: | Breeding | Migr. | Winter |
| <u>Buteo lagopus</u> | W W W W | W W W | | | | GL, Ag, S |
| | W W W W | W W W | | | | |
| | W W W W | W W W | | | | |
| | | | Abundance: | | C | |
| <u>Golden Eagle</u> | R R R R | R R R | Habitat: | Breeding GL, PJ, PP, | Migr. All | Winter All |
| <u>Aquila chrysaetos</u> | R R R R | R R R | | R, Ag, S, | | |
| | R R R R | R R R | | cliffs | | |
| | R R R R | R R R | | | | |
| | | | Abundance: | FC | FC | FC |
| <u>American Kestrel</u> | R R B R | R R R | Habitat: | Breeding Ag, RpL, S, | Migr. Ag, R, S, | Winter GL, Ag, RpL, |
| <u>Falco sparverius</u> | R R B R | R B R | | SgP, U, PP, | GL, U | S, U |
| | R B R R | R R R | | PJ, CG | | |
| | R R R R | B R R | | | | |
| | | | Abundance: | FC | FC-Ab | FC |
| <u>Merlin</u> | M M W | W W W | Habitat: | Breeding | Migr. | Winter |
| <u>Falco columbarius</u> | W M M W | W W W | | GL, RpL | GL, Ms, S, | |
| Acc. in 9, 21 | M W W W | W W M | | | R, Ag, U | |
| | W W W | W W W | | | | |
| | | | Abundance: | U | U | |
| <u>Peregrine Falcon</u> | B b b B | M M M | Habitat: | Breeding cliffs, CF, | Migr. GL, R, S, Aq | Winter R, Ms, GL, Sa |
| <u>Falco peregrinus</u> | R B b B | B M | | RpT, RpH | | |
| | B B B B | B M | | | | |
| | B B R b | B M M | | | | |
| | | | Abundance: | R | R | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|-------------------------------|-----------------|------------|--|------------------|
| <u>Gyrfalcon*</u> | | Breeding | Migr. | Winter |
| <u>Falco rusticolus</u> | M M | Habitat: | SgP, Ag | |
| Acc. in 5, 6, 11, 12, 18 | — — M M — — | | | |
| All records Dec. to Mar. | — — — — — — | Abundance: | R | |
| <u>Prairie Falcon</u> | | Breeding | Migr. | Winter |
| <u>Falco mexicanus</u> | R R R R R R R R | Habitat: | cliffs, R, GL, S, Ag, CF, GL, S, Cr, A | SgP, MmP, DS, Cr |
| | — — — — — — — — | | A | |
| | | Abundance: | U | FC |
| <u>Chukar</u> | | Breeding | Migr. | Winter |
| <u>Alectoris chukar</u> | R M | Habitat: | MmP, DS, Cr, PJ, SgSD, SaRb, | MmP, DS, Cr, Ag, |
| (Introduced) | — — b M — — | | barren | barren |
| Acc. in 4, 11, 12 | — — R — — — — | | ground | ground |
| | | Abundance: | U | U |
| <u>Ring-necked Pheasant</u> | | Breeding | Migr. | Winter |
| <u>Phasianus colchicus</u> | R R b R R R R | Habitat: | Ag, Cr, RPL | Ag, Cr, RPL |
| | — — — R R R R | | Ms | |
| (Introduced) | — — R R R R R R | | | |
| | | Abundance: | FC | FC |
| <u>Blue Grouse</u> | | Breeding | Migr. | Winter |
| <u>Dendragapus obscurus</u> | R R R R M — — | Habitat: | CF (except PJ), AT, Sc, MM, MmP, RpT, As | SF, DF, PP, LP |
| Acc. in 12 | — R R R R R b — | | | |
| | | Abundance: | FC | FC |
| <u>White-tailed Ptarmigan</u> | | Breeding | Migr. | Winter |
| <u>Lagopus leucurus</u> | R R R — — — — | Habitat: | A | AT |
| | — R R R — — — — | | | |
| | — R R R — — — — | | | |
| | — R R R — — — — | | | |
| | | Abundance: | FC | FC |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|---|------------------|----------|--|-------|--|
| <u>Northern Bobwhite</u> <u><i>Colinus virginianus</i></u> | R R R R | Habitat: | Breeding Ag,RpL | Migr. | Winter Ag,RpL |
| | — — — — | | | | |
| | R R R R | | | | |
| | — — — — | | | | |
| | R R R R | | | | |
| | — — — — | | | | |
| | R R R R | | | | |
| | — — — — | | | | |
| | Abundance: FC | | | FC | |
| <u>Scaled Quail</u> <u><i>Callipepla squamata</i></u> | M B — | Habitat: | Breeding SgSD,CG, SS,DS | Migr. | Winter Ag,RpL, RpT,DS,SS, CG,SgSD |
| Acc. in 5, 23 | — — — — | | | | |
| | R R R R R | | | | |
| | B b b R R R | | | | |
| | — — — — — | | | | |
| | Abundance: FC | | | FC | |
| <u>Gambel's Quail</u> <u><i>Callipepla gambelii</i></u> | — — — — — | Habitat: | Breeding MM,Sa,DS, SgP,Cr, barren areas | Migr. | Winter MM,Sa,DS, Cr,SgP, barren areas |
| | R R — — — | | | | |
| | R R — — — | | | | |
| | R — — — — | | | | |
| | — — — — — | | | | |
| | Abundance: FC | | | FC | |
| <u>Yellow Rail*</u> <u><i>Coturnicops noveboracensis</i></u> | — — — — — | Habitat: | Breeding | Migr. | Winter |
| Acc. in 21 | — — — — — | | | | |
| | — — — — M | | | | |
| | — — — — — | | | | |
| | Abundance: _____ | | | | |
| <u>Black Rail*</u> <u><i>Laterallus jamaicensis</i></u> | M — — — | Habitat: | Breeding | Migr. | Winter |
| Acc. in 4, 20 | — — — — — | | | Ms, | |
| | — — — — M — | | | | |
| | — — — — — | | | | |
| | Abundance: Und | | | | |
| <u>King Rail*</u> <u><i>Rallus elegans</i></u> | M — — — | Habitat: | Breeding | Migr. | Winter |
| Acc. in 5, 19 | — — — — — | | | Ms, | |
| | M — — — | | | | |
| | — — — — — | | | | |
| | Abundance: Und | | | | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | |
|--|--|------------|--------------------------------|
| <u>Virginia Rail</u> <u>Rallus limicola</u> | B B B R R R B R B B R R B R B R B b R B B B B B B b B b | Habitat: | Breeding Migr. Winter |
| | | Abundance: | FC C R |
| <u>Sora</u> <u>Porzana carolina</u> | B B B B B N B B B B B B B B b B b b B b b B B B B b B b | Habitat: | Breeding Migr. Winter |
| 6: One wintering record | | | Ms Ms |
| <u>Purple Gallinule*</u> <u>Porphyrrula martinica</u> | | Abundance: | FC C |
| Acc. in 23 | | Habitat: | Breeding Migr. Winter |
| | | | Ms, 1 record only |
| <u>Common Moorhen*/#</u> <u>Gallinula chloropus</u> | M M M M - - - - - - - - - - - - - - - | Habitat: | Breeding Migr. Winter |
| Acc. in 2, 4, 6, 8 | | | Ms, L |
| <u>American Coot</u> <u>Fulica americana</u> | B B B R R B B B B B R R R R B B B b R R B B R B B B b M | Habitat: | Breeding Migr. Winter |
| | | | Ms Ms, L Ms, L |
| | Abundance: | Ab | FC |
| <u>Sandhill Crane</u> <u>Grus canadensis</u> | M B B M M M M M M M M M M M M M M M M M M M M M M M M M | Habitat: | Breeding Migr. Winter |
| | | | MmP, Ms Ri, Cr, Sa, W/OG |
| | Abundance: | R-U | U-Ab |

STATUS
 R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE
 Ab - Abundant R - Rare
 C - Common Und - Undetermined
 FC - Fairly Common Irr - Irregular
 U - Unusual

| | | | | | |
|--|--|------------------------------------|-------------------|---------------|--------|
| <u>Whooping Crane</u> | <u>Grus americana</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 2, 3, 4, 5, 12, 15, 18, 19 Most records are birds from Grays Lake, Idaho population | M M M M M — M — M M — M — M M — | M — M — M — — — | | Ms,Cr,Sa | |
| | | <u>Abundance:</u> | R-U | | |
| <u>Black-bellied Plover</u> | <u>Pluvialis squatarola</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 1 | M — M M M — M M M — M — M M M | M M M M M M M M M M M M M | | W/OG,L, Cr | |
| | | <u>Abundance:</u> | R-U | | |
| <u>Lesser Golden Plover</u> | <u>Pluvialis dominica</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 6, 7, 8, 11, 13, 14, 16, 19, 21, 24, 25 Most records in the fall | M — M M M — — M M | M M M M M M — — — | | W/OG,L, Cr | |
| | | <u>Abundance:</u> | R | | |
| <u>Snowy Plover</u> | <u>Charadrius alexandrinus</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 5, 7, 12, 17, 24 25: No longer B-habitat gone | M — M M M — M B — M B | M — M M B B M — M | alkali flats | W/OG,L | |
| | | <u>Abundance:</u> | R, (C in 20) R | | |
| <u>Semipalmated Plover</u> | <u>Charadrius semipalmatus</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 10 | M — M M M — M M M M M M M M M M | M M M M M M M M M M M M | | W/OG,L | |
| | | <u>Abundance:</u> | U-FC | | |
| <u>Piping Plover</u> | <u>Charadrius melanotos</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 4, 11, 12, 14, 19, 21 | — — — M — — — M — — — M M M — — — — — | M — M M — M M M M — — — | | W/OG,L | |
| | | <u>Abundance:</u> | R | | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|--|--|--|-----------------|---|
| <u>Killdeer</u> <u>Charadrius vociferus</u> | R R B R R B B R R R R B B R B R | R R R R R G R R R R B B B B B | Habitat: | Breeding GL,Cr,Aq,U Migr. U Winter |
| | | | Abundance: C | C U-FC |
| <u>Mountain Plover</u> <u>Charadrius montanus</u> | b — M B M — B B M — B — B M | B B M B B M B B B B B B | Habitat: | Breeding SgP,MmP Migr. SgP,MmP, Cr,W/OG, L |
| Acc. in 7, 8, 14, 15, 25 | | | Abundance: R-FC | R-Ab |
| <u>Black-necked Stilt</u> <u>Himantopus mexicanus</u> | M M B M M M M M M M M M M B B | B — M M — M B B M M M M | Habitat: | Breeding alkali flats, Migr. W/OG,Ms, L,Ms |
| Acc. in 7, 10, 22 | | | Abundance: U | U |
| <u>American Avocet</u> <u>Recurvirostra americana</u> | B b B B B M M B B M M B M M B B | B B b B b M B B b B M M | Habitat: | Breeding L,Ms, alkali flats |
| | | | Abundance: FC | FC-Ab |
| <u>Greater Yellowlegs</u> <u>Tringa melanoleuca</u> | M M M M M M M M M M M M M M M M | M M M M M M M M M M M M | Habitat: | Breeding Ms,L,W/OG, Migr. Ri |
| Acc. in 10 | | | Abundance: FC | FC |
| <u>Lesser Yellowlegs</u> <u>Tringa flavipes</u> | M M n n M M M M M M M M M M M M | M M M M M M M M M M M M | Habitat: | Breeding L,Ri,Ms, Migr. W/OG |
| Acc. in 2 | | | Abundance: C | FC-C |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

Ab - Abundant R - Rare
C - Common Und - Undetermined
FC - Fairly Common Irr - Irregular
U - Unusual

| | | | | |
|------------------------------------|-------------------|------------|--------------|--------------|
| Solitary Sandpiper | | Breeding | Migr. | Winter |
| <u>Tringa solitaria</u> | M M M M M M M M | Habitat: | Aq | |
| Acc. in 9 | — M M M M M M M M | | | |
| | | Abundance: | FC | |
| Willet | | Breeding | Migr. | Winter |
| <u>Catoptrophorus semipalmatus</u> | b b B b M M M M | Habitat: | Ms | Ms,L,W/OG |
| Acc. in 9 | — M M M b M M M M | | | |
| | | Abundance: | Und | FC |
| Spotted Sandpiper | | Breeding | Migr. | Winter |
| <u>Actitis macularia</u> | B B B B B B b B | Habitat: | Aq | Aq |
| | — B B B B B B B B | | | |
| | | Abundance: | FC | FC |
| Upland Sandpiper | | Breeding | Migr. | Winter |
| <u>Bartramia longicauda</u> | M — M M B B B B | Habitat: | TgP,SgP | Cr,TgP,SgP |
| Acc. in 3, 4, 8, 11 | — — M M M M M M | | | |
| | | Abundance: | R | R |
| Whimbrel | | Breeding | Migr. | Winter |
| <u>Numenius phaeopus</u> | M — M M M M M M | Habitat: | L,Ms,W/OG | |
| Acc. in 7, 13, 20, 23 | — M M M M M M M | | | |
| | | Abundance: | R | |
| Long-billed Curlew | | Breeding | Migr. | Winter |
| <u>Numenius americanus</u> | b M b b B B b B | Habitat: | SgP,Cr-wheat | GL,Ms,L,W/OG |
| Acc. in 16 | — M M M M B B B B | | | |
| | | Abundance: | R-U | U |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|---|-----------------|------------|---------------|--------|
| Hudsonian Godwit | | Breeding | Migr. | Winter |
| <u>Limosa haemastica</u> | M M M M M | Habitat: | L,W/OG | |
| Acc. in 4, 5, 6, 8, 12, 19, 20, 21 | ----- | | | |
| | | Abundance: | R | |
| Marbled Godwit | | Breeding | Migr. | Winter |
| <u>Limosa fedoa</u> | M M M M B M M | Habitat: | L,W/OG, Ms | |
| Acc. in 9, 10, 14, 28 5: One breeding record, 1984 | M M M M M M M M | | | |
| | | Abundance: | FC | |
| Ruddy Turnstone | | Breeding | Migr. | Winter |
| <u>Arenaria interpres</u> | M M M M M M M M | Habitat: | W/OG,L | |
| Acc. in 1, 5, 6, 7, 8, 9, 11, 13, 14, 19, 20, 26 | ----- | | | |
| | | Abundance: | R | |
| Red Knot | | Breeding | Migr. | Winter |
| <u>Calidris canutus</u> | M M M M M M M M | Habitat: | L,W/OG | |
| Acc. in 5, 6, 7, 8, 14, 19, 21 | ----- | | | |
| | | Abundance: | R | |
| Sanderling | | Breeding | Migr. | Winter |
| <u>Calidris alba</u> | M M M M M M M M | Habitat: | W/OG,L,Ms | |
| Acc. in 8, 13 | ----- | | | |
| | | Abundance: | U | |
| Semipalmated Sandpiper | | Breeding | Migr. | Winter |
| <u>Calidris pusilla</u> | M M M M M M M M | Habitat: | L,Ms,W/OG | |
| Acc. in 2, 10, 13 | ----- | | | |
| | | Abundance: | FC | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

Western SandpiperCalidris mauri

Acc. in 10

| | | | | | | |
|---|---|---|---|---|---|---|
| M | M | M | M | M | M | M |
| M | M | M | M | M | M | M |
| M | M | M | M | M | M | M |
| M | M | M | M | M | M | M |

Habitat: Breeding Migr. Winter

L,Ms,Cr,
W/OGLeast SandpiperCalidris minutilla

Acc. in 9

| | | | | | | |
|---|---|---|---|---|---|---|
| M | M | M | M | M | M | M |
| M | M | - | M | M | M | M |
| M | M | M | M | M | M | M |
| M | M | M | M | M | M | M |

Habitat: Breeding Migr. Winter

L,Ms,W/OG

White-rumped SandpiperCalidris fuscicollis

Acc. in 8, 13, 16, 21

| | | | | | | |
|---|---|---|---|---|---|---|
| M | - | M | M | M | M | M |
| M | - | M | M | M | M | M |
| M | - | M | M | M | M | M |
| M | M | - | M | M | M | M |

Abundance: C

Habitat: Breeding Migr. Winter

L,W/OG

Baird's SandpiperCalidris bairdii

Acc. in 2

| | | | | | | |
|---|---|---|---|---|---|---|
| M | M | M | M | M | M | M |
| M | M | M | M | M | M | M |
| M | M | M | M | M | M | M |
| M | M | M | M | M | M | M |

Abundance: R

Habitat: Breeding Migr. Winter

L,W/OG

Pectoral SandpiperCalidris melanotos

| | | | | | | |
|---|---|---|---|---|---|---|
| M | M | M | M | M | M | M |
| M | M | M | M | M | M | M |
| M | M | M | M | M | M | M |
| - | - | M | M | M | M | M |

Abundance: FC-Ab

Habitat: Breeding Migr. Winter

L,W/OG

Sharp-tailed Sandpiper*Calidris acuminata

Acc. in 4

| | | | | | | |
|---|---|---|---|---|---|---|
| - | - | M | - | - | - | - |
| - | - | - | - | - | - | - |
| - | - | - | - | - | - | - |
| - | - | - | - | - | - | - |

Abundance: U

Habitat: Breeding Migr. Winter

L,W/OG

STATUS

- R-Resident (breeds)
- N-Nonbreeder (resident)
- B-Definite breeder
- b-Likely breeder
- n-Non-breeder (summer)
- W-Winter visitor
- M-Migrant

ABUNDANCE

- | | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|---|----------------------------|------------|-----------------------|--------|
| Dunlin | | Breeding | Migr. | Winter |
| <u>Calidris alpina</u> | M — — M M M | Habitat: | L,Ms,W/OG | |
| Acc. in 4, 8, 11, 21, 24, 25 | M M — M M M — M M — — — | | | |
| | | Abundance: | R | |
| Stilt Sandpiper | | Breeding | Migr. | Winter |
| <u>Calidris himantopus</u> | M — — M M M | Habitat: | L,Ms,W/OG | |
| Acc. in 15 | M M — M M M — M M — — — | | | |
| | | Abundance: | U-FC | |
| Buff-breasted Sandpiper | | Breeding | Migr. | Winter |
| <u>Tryngites subruficollis</u> | — — — M M M | Habitat: | L,Cr,SgP, TgP,W/OG | |
| Acc. in 5, 6, 7, 12, 14, 20, 21 | — — — M M M — — — — — — | | | |
| | | Abundance: | R | |
| Ruff*# | | Breeding | Migr. | Winter |
| <u>Philomachus pugnax</u> | — — — M — — | Habitat: | W/OG,L, 1 record | |
| Acc. in 5 | — — — — — — | | | |
| | | Abundance: | Und | |
| Short-billed Dowitcher* | | Breeding | Migr. | Winter |
| <u>Limnodromus griseus</u> | — — — M M M | Habitat: | L,Ms,W/OG, Cr | |
| Acc. in 6, 7, 11, 15, 16, 18, 19, 20 | M M — M M M | | | |
| | | Abundance: | R | |
| Long-billed Dowitcher | | Breeding | Migr. | Winter |
| <u>Limnodromus scolopaceus</u> | M M M M M M M | Habitat: | L,Ms,W/OG, Cr | |
| Acc. in 2,10 | M M M M M M M | | | |
| | | Abundance: | FC | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|--|------------------|--------------------|------------------------------|-----------|--------------------------|
| <u>Common Snipe</u> <u>Gallinago gallinago</u> | B B B R R R N | Habitat: | Breeding GL,R,Ms, W/OG | Migr. | Winter R,Aq RpL,Aq |
| | R B R R R R R | | | | |
| | R R R b b R R | | | | |
| | B R R R b B M | | | | |
| | | Abundance: | FC | FC | FC |
| <u>American Woodcock*</u> <u>Scolopax minor</u> | | Habitat: | Breeding | Migr. | Winter |
| Acc. in 4, 19 | ----- M ----- | | | Ms | |
| | ----- M ----- | | | | |
| | ----- ----- | | | | |
| <u>Wilson's Phalarope</u> <u>Phalaropus tricolor</u> | | Abundance: | R | | |
| | b B B B B b B | Habitat: | Breeding MmP,W/OG, | Migr. | Winter W/OG,Ms, |
| | b M M B B M M | | Ms,Cr | Cr,MmP | |
| | b B B B b B b | | | | |
| | b M B B b b M | | | | |
| <u>Red-necked Phalarope</u> <u>Phalaropus lobatus</u> | | Abundance: | FC | C-Ab | |
| Acc. in 9, 10, 21, 28 | M M M M M M M | Habitat: | Breeding | Migr. | Winter |
| | M M M M M M M | | | W/OG,Ms,L | |
| | M M M M M M M | | | | |
| | ----- M ----- | | | | |
| | | Abundance: | U-FC | | |
| <u>Red Phalarope</u> <u>Phalaropus fulicaria</u> | M M M M M M M | Habitat: | Breeding | Migr. | Winter |
| Acc. in 1, 4, 5, 6, 7, 8, 11, 12, 19, 24 | M M M M M M M | | | L | |
| | ----- M ----- | | | | |
| | ----- ----- | | | | |
| <u>Pomarine Jaeger</u> <u>Stercorarius pomarinus</u> | | Abundance: | R | | |
| Acc. in 5, 6, 12, 18 | ----- M M ----- | Habitat: | Breeding | Migr. | Winter |
| | ----- M ----- | | | L | |
| | ----- ----- | | | | |
| | | Abundance: | R | | |
| <u>STATUS</u> | | | | | |
| R-Resident (breeds) | | | | | |
| N-Nonbreeder (resident) | | | | | |
| B-Definite breeder | | | | | |
| b-Likely breeder | | | | | |
| n-Non-breeder (summer) | | | | | |
| W-Winter visitor | | | | | |
| M-Migrant | | | | | |
| | | <u>ABUNDANCE</u> | | | |
| Ab - Abundant | | R - Rare | | | |
| C - Common | | Und - Undetermined | | | |
| FC - Fairly Common | | Irr - Irregular | | | |
| U - Unusual | | | | | |

| | | | | |
|------------------------------------|-----------------|-----------------|----------------------|---------------|
| <u>Parasitic Jaeger</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Stercorarius parasiticus</u> | — M M M M | Habitat: | L | |
| Acc. in 5, 7, 11, 16, 19, 20 | — M — M M — | | | |
| | — — — — | | | |
| | | Abundance: | R | |
| <u>Laughing Gull#</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Larus atricilla</u> | — M M M — | Habitat: | L | |
| Acc. in 4, 5, 7, 11, 12, 19, 21 | — — — M M M | | | |
| | — — — — | | | |
| | | Abundance: | R | |
| <u>Franklin's Gull</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Larus pipixcan</u> | M M M M M M M M | Habitat: | Cr, Aq, GL | |
| Acc. in 2, 13 | M M M M M M M M | | | |
| | M M M M M M M M | | | |
| | | Abundance: | FC spring Ab fall | |
| <u>Little Gull*</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Larus minutus</u> | — M — M | Habitat: | L | |
| Acc. in 4, 7, 12 | — — — — | | | |
| | — — — — | | | |
| | | Abundance: | R | |
| <u>Bonaparte's Gull</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Larus philadelphia</u> | M M M M M M M M | Habitat: | L, Ms | |
| Acc. in 2, 9, 10, 13, 17, 21 | M M M M M M M M | | | |
| | M M M M M — — | | | |
| | | Abundance: | U-FC | |
| <u>Mew Gull*</u> | | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| <u>Larus canus</u> | — M — — | Habitat: | L | |
| Acc. in 4, 11, 12 | — — — — | | | |
| | — — — — | | | |
| | | Abundance: | Und | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|---|---------------------------|-------------------|--------------------------------|------------------|------------------|
| <u>Ring-billed Gull</u> | <u>Larus delawarensis</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| | | | L | L,Cr, U-dumps | L,Ri, U-dumps |
| | | | | | |
| | | | | | |
| | | | | | |
| <u>California Gull</u> | <u>Larus californicus</u> | <u>Abundance:</u> | R | FC-Ab | FC-Ab |
| | | <u>Habitat:</u> | Breeding | Migr. | Winter |
| | | | L | L,Cr, U-dumps | L,Ri, U-dumps |
| Acc. in 9 | | | | | |
| | | | | | |
| | | <u>Abundance:</u> | FC | FC | FC |
| <u>Herring Gull</u> | <u>Larus argentatus</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| | | | L | L, U-dumps | L,Ri, U-dumps |
| Acc. in 16, 22, 23, 25 | | | | | |
| | | | | | |
| | | <u>Abundance:</u> | R | FC | FC |
| <u>Thayer's Gull</u> | <u>Larus thayeri</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| | | | L, | L,Ri, U-dumps | L,Ri, U-dumps |
| Acc. in 14, 15, 20 | | | | | |
| | | | | | |
| | | <u>Abundance:</u> | R | R | R |
| <u>Lesser Black-backed Gull*</u> | <u>Larus fuscus</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| | | | L | | |
| Acc. in 11 | | | | | |
| | | | | | |
| | | <u>Abundance:</u> | Und | | |
| <u>Glaucous-winged Gull*</u> | <u>Larus glaucescens</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| | | | L, 1 record, Antero Res. | | |
| Acc. in 11, 18 (One record, same bird) | | | | | |
| | | | | | |
| | | <u>Abundance:</u> | Und | | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|--|-------------------------------------|------------|----------|--------|--------|
| <u>Glaucous Gull</u> <u>Larus hyperboreus</u> | | Habitat: | Breeding | Migr. | Winter |
| | W W W M | | L | | L |
| | W W W W | | | | |
| | W W W | | | | |
| | ----- | | | | |
| <u>Great Black-backed Gull*</u> <u>Larus marinus</u> | | Abundance: | R | U | |
| | | Breeding | Migr. | Winter | |
| | M | Habitat: | | | L |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| Acc. in 11 | | Abundance: | | R | |
| <u>Black-legged Kittiwake</u> <u>Rissa tridactyla</u> | | Habitat: | Breeding | Migr. | Winter |
| Acc. in 6, 8, 11, 19, 22 | M M M M M M M M M | | L | | |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| Acc. in 6, 8, 11, 19, 22 | Abundance: | | R | | |
| <u>Ross' Gull*</u> <u>Rhodostethia rosea</u> | | Habitat: | Breeding | Migr. | Winter |
| Acc. in 7 | M | | L | | |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| Acc. in 7 | Abundance: | | R | | |
| <u>Sabine's Gull</u> <u>Xema sabini</u> | | Habitat: | Breeding | Migr. | Winter |
| Acc. in 3, 6, 7, 8, 10, 14, 16, 18, 20, 24 | M M M M M M M M M M M M M M M | | L | | |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| Acc. in 3, 6, 7, 8, 10, 14, 16, 18, 20, 24 | Abundance: | | R | | |
| <u>Caspian Tern</u> <u>Sterna caspia</u> | | Habitat: | Breeding | Migr. | Winter |
| Acc. in 7, 8, 10, 11, 15, 16, 18, 19 | M M M M M M M M M M | | L, Ri | | |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| | ----- | | | | |
| Acc. in 7, 8, 10, 11, 15, 16, 18, 19 | Abundance: | | R | | |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|--|--|------------|--------------------------|--------|
| Common Tern | | Breeding | Migr. | Winter |
| <u>Sterna hirundo</u> | M M M M M M | Habitat: | L,Ms | |
| Acc. in 8, 14, 15, 16, 17, 18, 21 | M M M M M M M M M M M M M M M M M M | | | |
| | ----- | Abundance: | R-U | |
| Arctic Tern* | | Breeding | Migr. | Winter |
| <u>Sterna paradisaea</u> | M ----- | Habitat: | L | |
| Acc. in 4 | ----- | | | |
| | ----- | Abundance: | Und | |
| Forster's Tern | | Breeding | Migr. | Winter |
| <u>Sterna forsteri</u> | M B b B M b M M M B B M M b M n B B b M M b M b M | Habitat: | Ms,L | L,Ms |
| Acc. in 9, 15 | ----- | | | |
| | ----- | Abundance: | U | U-FC |
| Least Tern | | Breeding | Migr. | Winter |
| <u>Sterna antillarum</u> | M M M M M M | Habitat: | L,W/OG | L,Ri |
| Acc. in 4, 5, 7, 11, 14, 15, 16, 19, 26 | M M M B B B M M M B B B M M M B B B | | | |
| | ----- | Abundance: | R | R |
| Black Tern | | Breeding | Migr. | Winter |
| <u>Chlidonias niger</u> | M M B B B b b M M M M B M b b n M M B B b M M M B b M M | Habitat: | Ms | L,Ms |
| | ----- | Abundance: | U | U-FC |
| Marbled Murrelet* | | Breeding | Migr. | Winter |
| <u>Brachyramphus marmoratus</u> | M ----- | Habitat: | Und, 1 record only | |
| Acc. in 10 | ----- | | | |
| | ----- | Abundance: | Und | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | |
|---|--|------------|--|
| Rock Dove <u>Columba livia</u> (Introduced) | R R R R R R R R R R R R R R R R R R R R R R R R R R R R | Habitat: | Breeding Migr. Winter U,Ag, cliffs, bridges, cliffs dwellings |
| | | Abundance: | C C |
| Band-tailed Pigeon <u>Columba fasciata</u> | B B B B -- B B B B B b -- B B B B B b M | Habitat: | Breeding Migr. Winter PP,Sc,SF, Cr,Ag PP,SC,Cr, Ag |
| Acc. in 28 | | Abundance: | FC FC |
| White-winged Dove <u>Zenaida asiatica</u> | M M M M -- --- M M --- --- M M --- --- M M --- | Habitat: | Breeding Migr. Winter GL,S |
| Acc. in 1, 3, 5, 11, 12, 20, 27 | | Abundance: | R |
| Mourning Dove <u>Zenaida macroura</u> | R R B R R R B R R R B R B R B B B R R B R B B B R R R R | Habitat: | Breeding Migr. Winter GL,S,Ag, Cr,RpL, RpT,U,PP Ag,GL,S, U,RpL,Rpt Ag,GL,S,U, RpL |
| Black-billed Cuckoo <u>Coccyzus erythrophthalmus</u> | -- M B M b B -- B B M M M M M B B B B B M B M | Habitat: | Breeding Migr. Winter RpL RpL |
| Acc. in 3, 5, 15, 17, 26 | | Abundance: | Ab Ab U |
| Yellow-billed Cuckoo <u>Coccyzus americanus</u> | M M B B B B B B B B B B M B M M B B B | Habitat: | Breeding Migr. Winter RpL RpT,RpL |
| Acc. in 2, 3, 17, 22, 24, 25 | | Abundance: | R |
| | | Abundance: | U |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

Greater Roadrunner

Geococcyx californianus

Acc. in 11, 21, 22, 23, 24

| Habitat: | Breeding | Migr. | Winter |
|----------|----------------------------|-------|----------------------------|
| | DS,PJ, SaRb,SgSD, CG | | DS,PJ, SaRb,SgSD, CG |
| --- | M | | |
| --- | R | R b M | |
| M M M R | R R R | | |

Groove-billed Ani*

Crotophaga sulcirostris

Acc. in 4, 14, 19

| Abundance: | FC | | |
|------------|----------|--------|--------|
| Habitat: | Breeding | Migr. | Winter |
| | | RpL,Ag | |
| --- | M | M | |
| --- | M | | |
| --- | | | |
| Abundance: | | R | |

Common Barn-Owl

Tyto alba

Acc. in 3, 15, 16, 22

| Habitat: | Breeding | Migr. | Winter |
|------------|---------------------------------|-------|--------|
| | cliffs, dwellings, Ag,RpL | | RpL,Ag |
| --- | M R | R B b | |
| R | b | R B B | |
| M M b b | R R R | | |
| M M b | b R R | | |
| Abundance: | U | R | |

Flammulated Owl

Otus flammeolus

| Habitat: | Breeding | Migr. | Winter |
|------------|----------|-------|--------|
| | PP,As | PP,As | |
| --- | b | M | |
| B B | B | | |
| B B B | b | | |
| B B b B | b | | |
| Abundance: | FC | R | |

Eastern Screech-Owl

Otus asio

Acc. in 26

| Habitat: | Breeding | Migr. | Winter |
|------------|----------|-------|----------|
| | RpL,Ag,U | | RpL,Ag,U |
| --- | R b R N | | |
| --- | R R R | | |
| --- | M | | |
| Abundance: | U | U-FC | |

Western Screech-Owl

Otus kennicottii

| Habitat: | Breeding | Migr. | Winter |
|------------|---------------------|-------|---------------------|
| | RpL,RpT, Ag,U,PJ | | RpL,RpT, Ag,U,PJ |
| --- | R R b | | |
| R | | | |
| R b | R b | | |
| R B | B B | | |
| Abundance: | FC | U-FC | |

STATUS

R-Resident (breeds)

N-Nonbreeder (resident)

B-Definite breeder

b-Likely breeder

n-Non-breeder (summer)

W-Winter visitor

M-Migrant

ABUNDANCE

Ab - Abundant

R - Rare

C - Common

Und - Undetermined

FC - Fairly Common

Irr - Irregular

U - Unusual

| | | | | | |
|--|--------------------|---|-----------------|--------|--------------|
| Great Horned Owl <u>Bubo virginianus</u> | R R R R R R R | Habitat: R, CF, Ag | Breeding | Migr. | Winter |
| | R R R R R R R | | | | R, CF, Ag |
| | R R R R R R R | | | | |
| | R R R R R R R | | | | |
| | | | Abundance: FC-C | | FC-C |
| Snowy Owl <u>Nyctea scandiaca</u> | M W M W M M | Habitat: | Breeding | Migr. | Winter |
| Acc. in 2, 4, 6, 7, 8, 17, 18, 19, 21, 28 | M — W W — W | | | | Ag, GL, L |
| 11, 12: One wintering record each, 1977 and 1971 respectively | — M M M — M | | | | |
| | — — — — — M | | | | |
| | | | Abundance: | Irr, R | |
| Northern Pygmy-Owl <u>Glaucidium gnoma</u> | M R W — | Habitat: CF, As | Breeding | Migr. | Winter |
| Acc. in 1 | R N N R W — | | | | CF, RpT, RpL |
| | R R W R W — | | | | |
| | R R b b b — | | | | |
| | | | Abundance: U | | U |
| Burrowing Owl <u>Athene cunicularia</u> | B B b B B B B | Habitat: GL, rodent burrows | Breeding | Migr. | Winter |
| | B — B B B B | | | | |
| | B B B B B B B | | | | |
| | B b B b b B B | | | | |
| | | | Abundance: U-FC | | U |
| Spotted Owl <u>Strix occidentalis</u> | — M M M — | Habitat: | Breeding | Migr. | Winter |
| Acc. in 10, 11, 12, | — — — — — | | | | CF |
| | B — — — — — | | | | |
| | | | Abundance: | R | |
| Long-eared Owl <u>Asio otus</u> | R R b R b B B | Habitat: RpT, Sa, PJ, RpL, As, RpL, RPT, SaRb, DF, PJ, Sa Sc | Breeding | Migr. | Winter |
| Acc. in 3, 10, 18 | R B M R R — R | | | | RpL, Ag, PJ |
| | R W B M R B B | | | | |
| | b B R B — B B | | | | |
| | | | Abundance: U | U | Irr-U |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

Ab - Abundant R - Rare
C - Common Und - Undetermined
FC - Fairly Common Irr - Irregular
U - Unusual

| | | | | | | |
|------------------------------|---------------------------------|--|---------------|-----------------------------|-------------------------|-----------------|
| <u>Short-eared Owl</u> | <u>Asio flammeus</u> | M M R b R N | Habitat: | Breeding GL,Ms,Ag | Migr. GL,Ms,Ag | Winter GL,Ms,Ag |
| Acc. in 9, 18 | | M M B W R W R M M W M M M B R R M M N | | | | |
| <u>Boreal Owl</u> | <u>Aegolius funereus</u> | b R | Abundance: U | U | U | |
| Acc. in 3, 8, 9, 17 | | M M — W — M — — N b — | Habitat: | Breeding SF | Migr. SF | Winter SF |
| <u>Northern Saw-whet Owl</u> | <u>Aegolius acadicus</u> | b B B W | Abundance: R | R | R | |
| Acc. in 1, 15, 16 | | R N B R R — W b B B B B W R R N b B W | Habitat: | Breeding RpT,PP | Migr. RpT,RpL, Ag,PP | Winter |
| <u>Lesser Nighthawk*</u> | <u>Chordeiles acutipennis</u> | — — — — | Abundance: U | U | | |
| Acc. in 16 | | M — — — — — — — | Habitat: | Breeding RpL | Migr. | Winter |
| <u>Common Nighthawk</u> | <u>Chordeiles minor</u> | — — — — | Abundance: R | | | |
| | | B B B B B B B B B B B B b b B B B B B B B B B b B B B B | Habitat: | Breeding GL,R,U,PP, PJ,S | Migr. GL,R,U,S | Winter |
| <u>Common Poorwill</u> | <u>Phalaenoptilus nuttallii</u> | b b b M M M | Abundance: FC | Ab | | |
| Acc. in 7, 10 | | b b B B B — M b b B B B b — b b B B B B b | Habitat: | Breeding PJ,PP,S | Migr. RpT,RpL, PJ,S | Winter |
| | | | Abundance: FC | R | | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

Whip-poor-will*

Caprimulgus vociferus

Acc. in 4, 19, 27

| Habitat: | Breeding | Migr. | Winter |
|---------------|----------|-------|--------|
| ----- M ----- | | Sc | |
| ----- M ----- | | | |
| ----- M ----- | | | |
| | | | |

Black Swift

Cypseloides niger

Acc. in 3, 6, 12, 19

| Abundance: | Und | Breeding | Migr. | Winter |
|------------|---------|----------|-------|--------|
| Habitat: | cliffs | L,Ri | | |
| M b b B | M - | | | |
| b B b b | M ----- | | | |
| b B b b | ----- | | | |

Chimney Swift

Chaetura pelasgica

Acc. in 15, 17

| Abundance: | R | Breeding | Migr. | Winter |
|-------------|---------|----------|-------|--------|
| Habitat: | U | U,cliffs | | |
| M ----- B | B b b b | | | |
| M ----- B | B B B B | | | |
| ----- b b b | ----- | | | |

White-throated Swift

Aeronautes saxatalis

Acc. in 15, 17

| Habitat: | Breeding | Migr. | Winter |
|----------|-----------|--------|--------|
| b B b B | b B - | cliffs | cliffs |
| B B B B | B ----- | | |
| B B B B | B ----- | | |
| B B B B | B B ----- | | |

Blue-throated Hummingbird*

Lampornis clemenciae

Acc. in 4, 8, 11, 16, 19, 23

| Habitat: | Breeding | Migr. | Winter |
|-----------|----------|-------|--------|
| M ----- n | PP | PP,U | |
| M ----- n | | | |
| M ----- | | | |
| | | | |

Magnificent Hummingbird

Eugenes fulgens

Acc. in 3, 4, 9, 10, 18, 23, 25

| Habitat: | Breeding | Migr. | Winter |
|---------------|----------|----------|--------|
| M M ----- | RpT | RpT,PP,U | |
| M M ----- | | | |
| M M B ----- | | | |
| M M b M ----- | | | |

Und, 2
Abundance: records R

STATUS

R-Resident (breeds)

N-Nonbreeder (resident)

B-Definite breeder

b-Likely breeder

n-Non-breeder (summer)

W-Winter visitor

M-Migrant

ABUNDANCE

Ab - Abundant

R - Rare

C - Common

Und - Undetermined

FC - Fairly Common

Irr - Irregular

U - Unusual

| | | | | | |
|----------------------------------|--|------------|-----------------------------------|------------------|--------|
| <u>Black-chinned Hummingbird</u> | B b b M | Habitat: | Breeding | Migr. | Winter |
| <u>Archilochus alexandri</u> | <u>B B B M</u> | | PJ,RpT, RpL,U,Sc | PJ,RpT, RpL,U | |
| Acc. in 4, 11 | <u>B b B</u> <u>B B M B</u> | | | | |
| | <u>b b b</u> | | | | |
| | | Abundance: | FC | FC | |
| <u>Anna's Hummingbird*</u> | | | | | |
| <u>Calypte anna</u> | <u>M</u> <u>M</u> <u>M</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 8, 16, 19, 23 | <u>M</u> <u>M</u> <u>M</u> | | RpT,U | | |
| | <u>M</u> <u>M</u> <u>M</u> | | | | |
| <u>Calliope Hummingbird</u> | | Abundance: | R | | |
| <u>Stellula calliope</u> | <u>M M M M</u> <u>M M M M</u> <u>M M M M</u> <u>M M M M</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 6, 15 | <u>M</u> <u>M</u> <u>M</u> | | R,PP,T,U, late July- Sept. | | |
| | <u>M</u> <u>M</u> <u>M</u> | | | | |
| | | Abundance: | U-FC | | |
| <u>Broad-tailed Hummingbird</u> | | | | | |
| <u>Selasphorus platycercus</u> | <u>B B B B</u> <u>B B B B</u> <u>B B B B</u> <u>B B B B</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 13, 14 | <u>M M M</u> <u>B M M</u> <u>B M M</u> <u>B B M</u> | | S,CF,MmP, U,As,R | S,R,CF, MmP,T | |
| | | | | | |
| | | Abundance: | FC-C | FC | |
| <u>Rufous Hummingbird</u> | | | | | |
| <u>Selasphorus rufus</u> | <u>M M M M</u> <u>M M M M</u> <u>M M M M</u> <u>M M M M</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 2, 6, 27 | <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | CF,R,MmP, T,late July-Sept. | | |
| | | | | | |
| | | Abundance: | FC | | |
| <u>Belted Kingfisher</u> | | | | | |
| <u>Ceryle alcyon</u> | <u>R R R R</u> <u>R R R R</u> <u>R R R R</u> <u>B R R b</u> | Habitat: | Breeding | Migr. | Winter |
| 24: One wintering record | <u>R R R</u> <u>R M R</u> <u>R R R</u> <u>B B R</u> | | L,RpT,RpL, Ri | Ri | |
| | | | | | |
| | | Abundance: | FC | FC | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|-----------------------------------|-------------------|-------------------------|--------------|---------------------|
| Lewis' Woodpecker | | Breeding | Migr. | Winter |
| <u>Melanerpes lewis</u> | M B b R N M M | Sc, U, RpL, RpT, PJ, Ag | RpL, RpT, PJ | RpL, RpT, U, PJ, Ag |
| Acc. in 7 | R R M R R M | | | PP |
| | B R M R R R B | | | |
| | R R B R R R R | | | |
| | | U-FC west | | |
| | | Abundance: R-FC east | R | U |
| Red-headed Woodpecker | | Breeding | Migr. | Winter |
| <u>Melanerpes erythrocephalus</u> | n M M B B B B | Ag, RpL, U | Ag, RpL, PJ | |
| Acc. in 2, 3, 8, 10, 15, 16, 23 | M M B B B B | | | |
| | M M B B B B | | | |
| | M B b B B | | | |
| | | R west | | |
| | | Abundance: FC-C east | U | |
| Red-bellied Woodpecker | | Breeding | Migr. | Winter |
| <u>Melanerpes carolinus</u> | M M M M M | RpL | RpL, U, Ag | RpL |
| Acc. in 12 | ----- M M R | | | |
| | ----- M M M | | | |
| | ----- M M M M | | | |
| | | Abundance: R | R | R |
| Yellow-bellied Sapsucker | | Breeding | Migr. | Winter |
| <u>Sphyrapicus varius</u> | M M M M M M | RpL, U | | |
| Acc. in 6, 8, 11, 14, 15 | M M M M ----- | | | |
| | ----- ----- | | | |
| | | Abundance: R | | |
| Red-naped Sapsucker | | Breeding | Migr. | Winter |
| <u>Sphyrapicus nuchalis</u> | M B B B M -- | As | CF, R, As | RpL |
| Acc. in 5 | b B B B n -- | | | |
| | B B B B B -- | | | |
| | B B B B B b -- | | | |
| | | Abundance: FC | U | R |
| Williamson's Sapsucker | | Breeding | Migr. | Winter |
| <u>Sphyrapicus thyroideus</u> | B B B B M M | CF, As | CF, As, Sc | |
| Acc. in 14 | B B B B B -- | | | |
| | B B B B B -- | | | |
| | | Abundance: U-FC | U | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|---|--|----------------------------------|-----------------|--------------------|--------------------|-----------------------|
| <u>Ladder-backed Woodpecker</u> <u>Picoides scalaris</u> | M | Habitat: | Breeding | Migr. | Winter | |
| Acc. in 4, 21 | — — — R — — — M — — — M | R M M R R R | RpL, S, CG | RpL | RpL, PJ, CG | |
| | | | Abundance: U | R | U | |
| <u>Downy Woodpecker</u> <u>Picoides pubescens</u> | b R R R R R R R R R R R R R R R | R R R R b R R R B B B R | Habitat: | Breeding | Migr. | Winter |
| | | | PJ | CF, R, S, U, As | CF, R, S, U, As | CF, S, U, RpT, RpL |
| <u>Hairy Woodpecker</u> <u>Picoides villosus</u> | M R R R R R R R R R R B R R N R | R R R R — R R b b R B B | Abundance: FC | FC | FC | |
| | | | Habitat: | Breeding | Migr. | Winter |
| | | | SF, As, R | CF, R, S, As, U | CF, R, S, U | |
| <u>Three-toed Woodpecker</u> <u>Picoides tridactylus</u> | M R R R b R R R b b R R B R | M — — B — — B — — | Abundance: FC | FC | FC | |
| Acc. in 1, 12 | | | Habitat: | Breeding | Migr. | Winter |
| | | | SF | SF, PP, DF | | |
| <u>Northern Flicker</u> <u>Colaptes auratus</u> | R R B R R R R R R R R R R R R R | R R R R R R R R R R R R | Abundance: U | U | | |
| | | | Habitat: | Breeding | Migr. | Winter |
| | | | CF, U, As, R | | R, PP, U, Sc | |
| <u>Olive-sided Flycatcher</u> <u>Contopus borealis</u> | M b B B B B B B b B B B B B B B | M M M M — M b M M b b M | Abundance: C | C | | |
| Acc. in 7 | | | Habitat: | Breeding | Migr. | Winter |
| | | | SF, As, RpT, PP | R, Ag | | |
| | | | | | | |
| | | | Abundance: FC | FC | | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

Western Wood-Pewee

Contopus sordidulus

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| b | B | B | B | b | B | b | b |
| b | B | B | B | B | b | b | b |
| B | B | B | B | B | B | b | b |
| B | B | B | B | B | M | M | M |

Habitat: Breeding Migr. Winter

| | | |
|----------|----------|--|
| CF, RpL, | CF, RpL, | |
| RpT, As | RpT, U | |

Abundance: FC-C C

Eastern Wood-Pewee#

Contopus virens

Acc. 4, 6, 7, 19, 20

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| — | — | M | — | M | M | — | — |
| — | — | — | — | M | M | — | — |
| — | — | — | — | — | — | — | — |

Habitat: Breeding Migr. Winter

| | | |
|----------|--|--|
| RpL, Ag, | | |
| RpT | | |

Abundance: R

Willow Flycatcher

Empidonax traillii

Acc. in 5, 6, 7, 22, 28

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| b | b | B | B | M | M | M | — |
| b | b | B | B | M | M | — | — |
| b | B | B | B | b | — | — | — |
| M | b | M | b | b | b | M | — |

Habitat: Breeding Migr. Winter

| | | |
|-----------|-----------|--|
| RpL, RpT, | RpL, RpT, | |
| As | As | |

Abundance: FC FC

Least Flycatcher

Empidonax minimus

Acc. in 17, 21

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| — | — | B | — | M | M | M | — |
| — | — | M | — | M | — | M | — |
| — | — | M | M | M | M | M | — |
| — | — | M | — | — | — | M | — |

Habitat: Breeding Migr. Winter

| | | |
|-----|--|--|
| RpL | | |
|-----|--|--|

Abundance: U

Hammond's Flycatcher

Empidonax hammondi

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| b | B | B | B | M | — | — | — |
| B | B | B | B | M | — | — | — |
| b | b | B | B | B | — | — | — |
| B | B | B | B | B | — | — | — |

Habitat: Breeding Migr. Winter

| | | |
|----|-------|--|
| CF | CF, R | |
|----|-------|--|

Abundance: FC FC

Dusky Flycatcher

Empidonax oberholseri

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| B | B | B | B | M | — | — | — |
| B | b | B | B | b | — | — | — |
| B | B | B | B | B | — | — | — |
| B | B | B | B | B | — | — | — |

Habitat: Breeding Migr. Winter

| | | |
|-------------|-----------|--|
| PP, Sc, As, | R, As, Sc | |
| RpT, RpTT | | |

Abundance: FC FC

STATUS

R-Resident (breeds)

N-Nonbreeder (resident)

B-Definite breeder

b-Likely breeder

n-Non-breeder (summer)

W-Winter visitor

M-Migrant

ABUNDANCE

Ab - Abundant

R - Rare

C - Common

Und - Undetermined

FC - Fairly Common

Irr - Irregular

U - Unusual

| | | | | | |
|--|---|------------|--|-------------------------|--------------|
| <u>Gray Flycatcher</u> <u>Empidonax wrightii</u> | B B B B B <u>b</u> M b b b B B b M B | Habitat: | Breeding PJ | Migr. PJ | Winter |
| | | Abundance: | R | | |
| <u>Western Flycatcher</u> <u>Empidonax difficilis</u> | b B B B B B B B b B B B b B B B | Habitat: | Breeding Sa,SaRb, As,PP,RpT, CF,(often in dwell- ings) | Migr. R,As | Winter |
| Acc. in 7, 14, 21 | M M M B _ M B M M B B M | | | | |
| | | Abundance: | FC | R | |
| <u>Black Phoebe*</u> <u>Sayornis nigricans</u> | ----- - M - - - M - - | Habitat: | Breeding Ri,bridges | Migr. RpT | Winter |
| Acc. in 16, 23 19: Nested 1972-74, 1987. | - - - - - - - - - - - - | | | | |
| | | Abundance: | R | | |
| <u>Eastern Phoebe</u> <u>Sayornis phoebe</u> | ----- - M - - - M - - - M M b b M - - - - B b | Habitat: | Breeding RpL,Aq 0-dirt bank | Migr. RpL,RpT,U | Winter |
| Acc. in 5, 14, 17, 21 | | | | | |
| | | Abundance: | R | | |
| <u>Say's Phoebe</u> <u>Sayornis saya</u> | B B b B R b B B B B B B B B B B | Habitat: | Breeding Ag,S,GL, U,bridges | Migr. RpL,RpT,Ag, GL | Winter |
| | B B M B B B B B b B B B | | | | |
| | | Abundance: | FC | | |
| <u>Vermilion Flycatcher</u> <u>Pyrocephalus rubinus</u> | ----- M - - M - M - - - M M M | Habitat: | Breeding Ag,RpL, 1 record | Migr. RpL,RpT | Winter GL,Ag |
| Acc. in 4, 5, 6, 8, 12, 17, 19, 20, 21, 23, 24, 25, 26 6: Nested, 1981 | M B - M M M M - - | | | | |
| | | Abundance: | Und | R | |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

Ab - Abundant R - Rare
C - Common Und - Undetermined
FC - Fairly Common Irr - Irregular
U - Unusual

| | | | | | | |
|----------------------------------|-------------------------------------|----------------------------|----------------------------|------------|--------------|------------|
| <u>Ash-throated Flycatcher</u> | B B M M | M M | Habitat: | Breeding | Migr. | Winter |
| <u>Myiarchus cinerascens</u> | <u>B</u> <u>M</u> | <u>M</u> <u>M</u> | | PJ, RpL, | PJ, RpL, | |
| Acc. in 3, 10, 14 | <u>B</u> <u>B</u> | <u>M</u> <u>B</u> | | RpT | RpT, Ag, | |
| 11: Nested, 1986 | <u>B</u> <u>B</u> | <u>b</u> | <u>B</u> <u>M</u> | | SaRb | |
| | <u>B</u> <u>B</u> | <u>b</u> | <u>B</u> <u>B</u> <u>B</u> | Abundance: | FC | R |
| <u>Great Crested Flycatcher</u> | | | | | | |
| <u>Myiarchus crinitus</u> | <u>M</u> | <u>M</u> <u>B</u> <u>M</u> | Habitat: | Breeding | Migr. | Winter |
| | <u>M</u> | <u>M</u> <u>M</u> | | RpL | RpL, Ag | |
| Accidental in 4, 5, 19, 21, | <u>M</u> <u>M</u> <u>M</u> | | | | | |
| 23, 27, 28 | <u>M</u> <u>M</u> | | | | | |
| | | | Abundance: | R | R | |
| <u>Cassin's Kingbird</u> | | | | | | |
| <u>Tyrannus vociferans</u> | <u>M</u> | <u>b</u> <u>B</u> | <u>B</u> <u>M</u> <u>M</u> | Habitat: | Breeding | Migr. |
| | <u>b</u> <u>b</u> <u>B</u> <u>B</u> | <u>b</u> <u>b</u> <u>B</u> | <u>b</u> <u>b</u> <u>B</u> | | Ag, RpL, PJ, | Ag, RpL, |
| Acc. in 1 | <u>b</u> <u>b</u> | <u>B</u> | <u>B</u> <u>B</u> <u>B</u> | | CG, DS | SS, CG, DS |
| | <u>B</u> <u>B</u> <u>b</u> | <u>B</u> | <u>B</u> <u>B</u> <u>B</u> | | | |
| | | | Abundance: | U-FC | U-FC | |
| <u>Western Kingbird</u> | | | | | | |
| <u>Tyrannus verticalis</u> | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>B</u> <u>B</u> | Habitat: | Breeding | Migr. | Winter |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>B</u> <u>B</u> | | Ag, RpL, | Ag, RpL, U | |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>B</u> <u>B</u> | | RpT, U, PJ | | |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>B</u> <u>B</u> | | | | |
| | | | FC west | | | |
| | | | Abundance: | C-Ab east | C | |
| <u>Eastern Kingbird</u> | | | | | | |
| <u>Tyrannus tyrannus</u> | <u>n</u> <u>B</u> <u>b</u> <u>B</u> | <u>B</u> <u>B</u> <u>B</u> | Habitat: | Breeding | Migr. | Winter |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>B</u> <u>B</u> | | Ag, RpL, | Ag, RpL, U | |
| | <u>B</u> <u>b</u> <u>b</u> <u>B</u> | <u>B</u> <u>B</u> <u>B</u> | | RpT, SaRb, | | |
| | <u>b</u> <u>B</u> <u>M</u> <u>b</u> | <u>B</u> <u>B</u> <u>B</u> | | U, PJ | | |
| | | | U west | | | |
| | | | Abundance: | FC east | FC | |
| <u>Scissor-tailed Flycatcher</u> | | | | | | |
| <u>Tyrannus forficatus</u> | <u>M</u> | <u>M</u> <u>M</u> <u>M</u> | Habitat: | Breeding | Migr. | Winter |
| | <u>M</u> <u>M</u> | <u>M</u> <u>M</u> <u>M</u> | | RpL | RpL | |
| Acc. in 5, 6, 7, 810, 11, 13, | <u>M</u> <u>M</u> <u>M</u> | <u>M</u> <u>M</u> <u>M</u> | | | | |
| 17, 18, 19, 21, 22, 23 | <u>M</u> <u>M</u> <u>M</u> | <u>M</u> <u>B</u> | | | | |
| | | | FC-28 | | | |
| | | | R- | | | |
| | | | Abundance: | elsewhere | R | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|--|-------------------------------------|----------------------------|------------|----------------------------|--------------------------|------------------------|
| Horned Lark | R R R R | R R R | Habitat: | Breeding GL,T,U, Sa,DS | Migr. GL,Sa | Winter GL,Sa, SaRb, |
| <u>Eremophila alpestris</u> | <u>R</u> <u>R</u> <u>R</u> <u>R</u> | <u>R</u> <u>R</u> <u>R</u> | | | | |
| | <u>R</u> <u>D</u> <u>D</u> <u>R</u> | <u>R</u> <u>R</u> <u>R</u> | | | | |
| | <u>R</u> <u>R</u> <u>B</u> <u>R</u> | <u>R</u> <u>R</u> <u>R</u> | | | | |
| | <u>R</u> <u>R</u> <u>R</u> <u>R</u> | <u>R</u> <u>R</u> <u>R</u> | | | | |
| | | | Abundance: | C-Ab | Ab | Ab |
| Purple Martin | M M M | M M M | Habitat: | Breeding PP,RpT,As | Migr. PP,RpT,RpL | Winter |
| <u>Progne subis</u> | <u>M</u> <u>B</u> <u>M</u> <u>M</u> | <u>M</u> <u>—</u> <u>M</u> | | | | |
| Acc. in 1, 2, 4, 6, 7, 10, 11, 12, 14, 16, 25, 27 | <u>B</u> <u>M</u> <u>—</u> <u>—</u> | <u>—</u> <u>—</u> <u>—</u> | | | | |
| | <u>B</u> <u>M</u> <u>—</u> <u>M</u> | <u>—</u> <u>M</u> <u>—</u> | | | | |
| | | | Abundance: | U west | R | |
| Tree Swallow | B B B B | B M B | Habitat: | Breeding CF,As,RpL | Migr. R,Aq | Winter |
| <u>Tachycineta bicolor</u> | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>M</u> <u>M</u> | | | | |
| | <u>b</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>M</u> <u>M</u> | | | | |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>M</u> <u>M</u> | | | | |
| | | | Abundance: | C | C-Ab | |
| Violet-green Swallow | B B B B | M M | Habitat: | Breeding CF,As, RpL,RpT | Migr. R,Aq | Winter |
| <u>Tachycineta thalassina</u> | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>—</u> <u>M</u> | | | | |
| Acc. in 7 | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>M</u> <u>—</u> | | | | |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>B</u> <u>—</u> | | | | |
| | | | Abundance: | C | C-Ab | |
| Northern Rough-winged Swallow | B B B B | B B B | Habitat: | Breeding banks-R,Ag | Migr. R,Ag,Aq, MmP,Sa | Winter |
| <u>Stelgidopteryx serripennis</u> | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>b</u> <u>B</u> | | | | |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>B</u> <u>b</u> | | | | |
| | <u>B</u> <u>B</u> <u>b</u> <u>b</u> | <u>B</u> <u>B</u> <u>B</u> | | | | |
| | | | Abundance: | FC | FC-C | |
| Bank Swallow | b B B B | B B b | Habitat: | Breeding banks-R,Ag | Migr. Ag,Aq | Winter |
| <u>Riparia riparia</u> | <u>B</u> <u>D</u> <u>M</u> <u>B</u> | <u>B</u> <u>—</u> <u>b</u> | | | | |
| | <u>b</u> <u>B</u> <u>M</u> <u>b</u> | <u>B</u> <u>b</u> <u>b</u> | | | | |
| | <u>B</u> <u>B</u> <u>b</u> <u>B</u> | <u>b</u> <u>b</u> <u>b</u> | | | | |
| | | | Abundance: | FC | U-FC | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|--|--|--|------------|---------------------------------------|--------------------|------------------------|
| <u>Cliff Swallow</u> <u>Hirundo pyrrhonota</u> | B B B B B B B B B B B B B B B B | B B B B B B B B B B B B B B B B | Habitat: | Breeding cliffs, structures-R, Ag, Aq | Migr. Ag, Aq | Winter |
| | | | Abundance: | Ab | Ab | |
| <u>Barn Swallow</u> <u>Hirundo rustica</u> | B B B B B B B B B B B B B B B B | B B B B B B B B B B B B B B B B | Habitat: | Breeding structures- R, Ag, Aq | Migr. R, Ag, Aq | Winter |
| | | | Abundance: | C | Ab | |
| <u>Gray Jay</u> <u>Perisoreus canadensis</u> | R R R R B R R M R R R D R R R | — — — — — — W R — — | Habitat: | Breeding SF, LP | Migr. CF, RpT | Winter |
| Acc. in 15 | | | Abundance: | FC | FC | |
| <u>Steller's Jay</u> <u>Cyanocitta stelleri</u> | R R R R R R R R R R R R R R R R | M — — R — — R W R M M | Habitat: | Breeding CF | Migr. CF, As, MM | Winter CF, As, MM |
| | | | Abundance: | FC-C | FC-C | FC west |
| | | | | | Irr east | Irr east |
| <u>Blue Jay</u> <u>Cyanocitta cristata</u> | M R M R M R M M | R R R R b R R R R N D R | Habitat: | Breeding RpL, U, Ag | Migr. RpL, RpT, Ag | Winter RpL, RpT, Ag, U |
| Acc. in 8, 10, 15, 23, 24 | | | Abundance: | FC | R | U-FC |
| <u>Scrub Jay</u> <u>Aphelocoma coerulescens</u> | R R b R R R R R R R b R R R b R | M — — R — — R W R R M | Habitat: | Breeding Sc, PJ, MM | Migr. Sc, PJ, MM, | Winter U |
| Acc. in 5, 28 | | | Abundance: | FC | FC | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|--|--|--|-----------------|--|-----------------------------------|
| <u>Pinyon Jay</u> <u>Gymnorhinus cyanocephalus</u> | R R R M M R N R N M — M R R R R R W R R R R R b B | Habitat: PJ | Breeding | Migr. | Winter |
| | | | | PJ, RpL, RpT, SgP, SaRb, U, Sc, T | PJ, RpL, RpT, SgP, SaRb, Sc |
| | | | FC west | FC west | FC west |
| | | Abundance: Irr east | | Irr east | Irr east |
| <u>Clark's Nutcracker</u> <u>Nucifraga columbiana</u> | R R R R M — M R R R R M — M R R R R R W — R R R R R M — | Habitat: AT, LP, SF, DF | Breeding | Migr. | Winter |
| | | | | CF, A | SF, DF |
| | | | Abundance: FC | Irr east | Irr east |
| <u>Black-billed Magpie</u> <u>Pica pica</u> | R | Habitat: Ag, R, U, S, CG, PJ | Breeding | Migr. | Winter |
| | | | | | Ag, RpL, CG, RpT, U, S |
| | | | Abundance: C-Ab | C-Ab | |
| <u>American Crow</u> <u>Corvus brachyrhynchos</u> | R R R R R B R R b R R R R R R R R R R R R R R R R R R R | Habitat: Ag, U, RpL, RpT, PP, LP | Breeding | Migr. | Winter |
| | | | | All | Ag, U, RpL, RpT, S |
| | | | Abundance: FC | C-Ab | C-Ab |
| <u>Chihuahuan Raven</u> <u>Corvus cryptoleucus</u> | — — — M M b B — M — M R R R b — — b R R | Habitat: struct- ures-SgP, SgSD, MXP, TgP, SaRb, CG, TgP, CS, SS | Breeding | Migr. | Winter |
| Acc. in 5, 12, 16 | | | | | SgP, SgSD, MXP, TgP, SaRb, CG, SS |
| | | | Abundance: U | R west | U |
| <u>Common Raven</u> <u>Corvus corax</u> | R R R R R W W R R R R N W W R R R R R M M R R R R R R b | Habitat: A, CF, cliffs | Breeding | Migr. | Winter |
| | | | | A, CF, R, GL, Ag | A, CF, R, GL, Ag |
| | | | Abundance: FC | FC west | FC west |
| | | | | Irr east | Irr east |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|--------------------------------------|--|--|------------------------|------------------------------|
| <u>Black-capped Chickadee</u> | | Breeding | Migr. | Winter |
| <u>Parus atricapillus</u> | W R R R R R b R R R R R b R R R R R R R R R R R R R R M | RpL, RpT, U, Ag | R, U, Ag, As, S | RpL, RpT, U, Ag, As, S |
| Acc. in 28 | | Habitat: | | |
| | | Abundance: | FC-C | FC-C |
| <u>Mountain Chickadee</u> | | Breeding | Migr. | Winter |
| <u>Parus gambeli</u> | R R R R R W R R R R R M — R R R R R W — R R R R R W M | CF | | CF, Sc, As, U, RpT |
| Acc. in 13, 28 | | Habitat: | | |
| | | Abundance: | C | FC-AB |
| <u>Plain Titmouse</u> | | Breeding | Migr. | Winter |
| <u>Parus inornatus</u> | B R M — — R b R — — R R b R R — R R B B R M | PJ | PJ, PP, RpL, Sc | PJ, PP, RpL, Sc |
| Acc. in 4 | | Habitat: | | |
| | | Abundance: | FC | FC |
| <u>Bushtit</u> | | Breeding | Migr. | Winter |
| <u>Psaltriparus minimus</u> | b M R R R b W W — — R R b R R W R R N R b R — | PJ, S, RpL, RpT, Ag- orchards, west | | PJ, S, U, RpL, Ag-west |
| 12: Flock of 20 wintered, 1987-88 | | Habitat: | | |
| | | Abundance: | U | U |
| <u>Red-breasted Nuthatch</u> | | Breeding | Migr. | Winter |
| <u>Sitta canadensis</u> | R R b R M M W R b R R R M M M B b R R W M — R N R B M M | CF, RpH, RpT | CF, R, U, RpL | CF, R, U |
| Acc. in 5, 13, 27, 28 | | Habitat: | | |
| | | Abundance: | U-FC | U-C |
| <u>White-breasted Nuthatch</u> | | Breeding | Migr. | Winter |
| <u>Sitta carolinensis</u> | R R R R W M M R N R R R — W R R R R R W R R R R R R W — | CF, As, Sc, RpT | RpL, RpT, As, U, CF | CF, U, RpL, RpT |
| | | Habitat: | | |
| | | Abundance: | FC | U-FC |
| | | | | FC |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

Pygmy Nuthatch

Sitta pygmaea

Acc. in 2, 5, 6

| | | | | |
|---|---|---|---|---|
| b | R | R | M | M |
| R | b | R | R | R |
| B | R | B | R | R |
| R | R | R | R | b |

| | | | | | |
|--|--|--|----------|-------|--------|
| | | | Breeding | Migr. | Winter |
| | | | PP,DF | R,PP | PP |

Abundance: FC

Irr

FC

Brown Creeper

Certhia americana

Acc. in 2

| | | | | | |
|---|---|---|---|---|---|
| M | R | R | W | W | W |
| R | b | R | R | R | W |
| b | b | R | R | R | W |
| R | R | R | R | M | W |

| | | | | | |
|--|--|--|----------|---------------|------------------|
| | | | Breeding | Migr. | Winter |
| | | | CF,R | As,CF,U, R | CF,RpT, RpL,U |

Abundance: U-FC

U-FC

FC

Rock Wren

Salpinctes obsoletus

Acc. in 21

| | | | | | | |
|---|---|---|---|---|---|---|
| B | B | b | B | B | M | M |
| R | b | B | B | b | B | B |
| R | B | B | R | R | B | M |
| B | B | B | B | B | B | B |

| | | | | | |
|--|--|--|------------------|--------------------------------|--------------------------------|
| | | | Breeding | Migr. | Winter |
| | | | rocks, cliffs | Sg,rocks, SgSD,Sc, MM,CG | Sg,rocks, SgSD,Sc, MM,CG |

Abundance: FC

U

U

Canyon Wren

Catherpes mexicanus

Acc. in 6, 9, 10

| | | | | |
|---|---|---|---|---|
| B | R | R | M | — |
| R | M | M | R | R |
| B | B | B | R | B |
| B | B | b | B | R |

| | | | | | |
|--|--|--|----------|--------|--------|
| | | | Breeding | Migr. | Winter |
| | | | cliffs | cliffs | cliffs |

Abundance: FC

FC

U

Carolina Wren

Thryothorus ludovicianus

Acc. in 11, 12, 14, 19

| | | | |
|---|---|---|---|
| — | — | M | — |
| — | — | M | M |
| — | — | M | — |
| — | — | — | — |

| | | | | | |
|--|--|--|----------|-------|--------|
| | | | Breeding | Migr. | Winter |
| | | | RpL,U | RpL,U | RpL,U |

Abundance: R

Bewick's Wren

Thryomanes bewickii

Acc. in 4, 5, 12

| | | | | | |
|---|---|---|---|---|---|
| b | B | — | M | M | — |
| R | b | — | M | M | — |
| R | R | M | R | R | — |
| b | R | — | M | B | R |

| | | | | | |
|--|--|--|------------|---------|----------------|
| | | | Breeding | Migr. | Winter |
| | | | PJ,RpL,RpT | RpL,RpT | PJ,RpL, RpT |

Abundance: U-FC

R

U

STATUS

R-Resident (breeds)

N-Nonbreeder (resident)

B-Definite breeder

b-Likely breeder

n-Non-breeder (summer)

W-Winter visitor

M-Migrant

ABUNDANCE

Ab - Abundant

R - Rare

C - Common

Und - Undetermined

FC - Fairly Common

Irr - Irregular

U - Unusual

| | | | | | | |
|--------------------------------|----------------|--------------|------------|----------------|-----------------------|-----------------------|
| <u>House Wren</u> | B B B B | B B B | Habitat: | Breeding | Migr. | Winter |
| <u>Troglodytes aedon</u> | <u>B B B B</u> | <u>B B B</u> | | As, CF, R, | R, S | |
| | <u>B B B B</u> | <u>B B B</u> | | Sc, Sa, | | |
| | <u>B B B B</u> | <u>B B b</u> | | SaRb, SgSD, | | |
| | <u>B B B B</u> | <u>B B B</u> | | AT | | |
| | | | Abundance: | C | C | |
| <u>Winter Wren</u> | | | | | | |
| <u>Troglodytes troglodytes</u> | <u>W</u> | <u>W</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 9, 10, 16, 17, 28 | <u>M M W</u> | <u>W W</u> | | | | RpL, RpT |
| | <u>M M</u> | <u>W</u> | | | | |
| | <u>---</u> | <u>M M</u> | | | | |
| | | | Abundance: | | R | |
| <u>Sedge Wren*</u> | | | | | | |
| <u>Cistothorus platensis</u> | <u>M</u> | <u>M</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. 4, 5, 7, 14, 19, 26 | <u>M</u> | <u>M</u> | | Ms | Ms | |
| | <u>---</u> | <u>M</u> | | | | |
| | <u>---</u> | <u>M</u> | | | | |
| | <u>---</u> | <u>M</u> | | | | |
| | | | Abundance: | Und | R | |
| <u>Marsh Wren</u> | | | | | | |
| <u>Cistothorus palustris</u> | <u>B B b R</u> | <u>B W W</u> | Habitat: | Breeding | Migr. | Winter |
| | <u>R W b B</u> | <u>W R</u> | | Ms | Ms | Ms |
| | <u>W R M W</u> | <u>b R b</u> | | | | |
| | <u>B B R B</u> | <u>M W W</u> | | | | |
| | | | Abundance: | U | U | U |
| <u>American Dipper</u> | | | | | | |
| <u>Cinclus mexicanus</u> | <u>B R R R</u> | <u>M</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 7 | <u>R R R R</u> | <u>R</u> | | Ri, bridges | | Ri |
| | <u>R R R R</u> | <u>R W</u> | | | | |
| | <u>R R R R</u> | <u>B</u> | | | | |
| | | | Abundance: | FC | | FC |
| <u>Golden-crowned Kinglet</u> | | | | | | |
| <u>Regulus satrapa</u> | <u>W R R</u> | <u>W W W</u> | Habitat: | Breeding | Migr. | Winter |
| | <u>R B B R</u> | <u>W W W</u> | | SF, Lm, PJ | CF, Sc, RpL | CF, RpL, RpT, U |
| | <u>R B R R</u> | <u>R W W</u> | | | | |
| | <u>W R B B</u> | <u>b</u> | | | | |
| | | | Abundance: | U-FC | FC mtns Irr plains | FC mtns Irr plains |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

Ruby-crowned Kinglet

Regulus calendula

| Breeding | Migr. | Winter |
|----------------|--------------|-----------------|
| CF,AT | R,CF | RpL,PJ,U, PP |
| <u>M R B B</u> | <u>M M M</u> | |
| <u>R B B B</u> | <u>M M M</u> | |
| <u>R B B B</u> | <u>B M M</u> | |
| <u>R R B B</u> | <u>B M M</u> | |

Abundance:FC

FC-C

U-FC

Blue-gray Gnatcatcher

Polioptila caerulea

Acc. in 7

| Breeding | Migr. | Winter |
|----------------|--------------|----------|
| PJ,Sc, | S,PJ,RpT, | |
| SaRb,RpL, | RpL,U | |
| MM | | |
| <u>B B b B</u> | <u>M M M</u> | Habitat: |
| <u>B b b B</u> | <u>b M M</u> | |
| <u>B B B B</u> | <u>B M —</u> | |
| <u>B B — B</u> | <u>M b b</u> | |

Abundance:FC

FC

Eastern Bluebird

Sialia sialis

Acc. in 9, 13, 16

| Breeding | Migr. | Winter |
|----------------|--------------|----------|
| RpL,As | RpL,Ag | |
| <u>— M M B</u> | <u>b b B</u> | Habitat: |
| <u>— M n M</u> | <u>M M B</u> | |
| <u>— M n M</u> | <u>n M b</u> | |
| <u>— M M M</u> | <u>M M M</u> | |

Western Bluebird

Sialia mexicana

Acc. in 5, 6, 14, 20, 28

| Breeding | Migr. | Winter |
|----------------|--------------|-----------|
| PP,Sc,PJ, | PP,GL,PJ, | PP,Sc,PJ, |
| U | Ag,MM,As, | RpL,RpT, |
| | RpL,RpT,U | U,SgSD |
| <u>B b B B</u> | <u>M M —</u> | Habitat: |
| <u>B M b B</u> | <u>B M —</u> | |
| <u>B R B B</u> | <u>B M —</u> | |
| <u>R R B B</u> | <u>B — M</u> | |

Mountain Bluebird

Sialia currucoides

| Breeding | Migr. | Winter |
|----------------|--------------|-----------|
| MM,As,PP, | Ag,U,GL | Ag,PJ,MM, |
| PJ,Sc | | SaRb |
| <u>B B B B</u> | <u>B M M</u> | Habitat: |
| <u>B B B B</u> | <u>B M —</u> | |
| <u>B R B R</u> | <u>B M —</u> | |
| <u>R R B B</u> | <u>R B M</u> | |

Abundance:U-FC

U

U-FC

Townsend's Solitaire

Myadestes townsendi

| Breeding | Migr. | Winter |
|----------------|--------------|------------|
| CF,As | CF,RpL, | PP,PJ,U, |
| | RpT | RpL,RpT,Ag |
| <u>M b R R</u> | <u>W W W</u> | Habitat: |
| <u>R B R R</u> | <u>R W W</u> | |
| <u>R R R R</u> | <u>R W W</u> | |
| <u>R R R R</u> | <u>b W W</u> | |

Abundance:FC

FC

FC-C

STATUS

R-Resident (breeds)

N-Nonbreeder (resident)

B-Definite breeder

b-Likely breeder

n-Non-breeder (summer)

W-Winter visitor

M-Migrant

ABUNDANCE

Ab - Abundant

R - Rare

C - Common

Und - Undetermined

FC - Fairly Common

Irr - Irregular

U - Unusual

| | | | | |
|-----------------------------|--|-------------------------------------|----------------------------|----------------------------|
| <u>Veery</u> | | Breeding | Migr. | Winter |
| <u>Catharus fuscescens</u> | B B b B B B B B B b M M M | Habitat: RpT, RPL | RpT, RPL | |
| Acc. in 26 | | | | |
| | | Abundance: U-FC | U | |
| <u>Gray-cheeked Thrush</u> | | Breeding | Migr. | Winter |
| <u>Catharus minimus</u> | M M M M M M M | Habitat: | U, Ag, RPL | |
| Acc. in 4, 5, 7, 13, 28 | M | | | |
| | | Abundance: R | | |
| <u>Swainson's Thrush</u> | | Breeding | Migr. | Winter |
| <u>Catharus ustulatus</u> | M b b B B B B B b B B B B B B B | Habitat: RpT, RPL, SF | R, U, Ag, Sc | |
| | M M M B M M B M M B M M | | | |
| | | Abundance: FC | FC-Ab | |
| <u>Hermit Thrush</u> | | Breeding | Migr. | Winter |
| <u>Catharus guttatus</u> | B B B B B B B B B B B B B B B B | Habitat: CF | CF, R, U, Ag | |
| | M M M B M M B M M B M M | | | |
| | | Abundance: FC | FC | |
| <u>Wood Thrush</u> | | Breeding | Migr. | Winter |
| <u>Hylocichla mustelina</u> | M M M M M M M M M M M M M M M M | Habitat: | RPL, RpT | |
| Acc. in 6, 8, 17, 23 | M M M M | | | |
| | | Abundance: R | | |
| <u>American Robin</u> | | Breeding | Migr. | Winter |
| <u>Turdus migratorius</u> | R R R R R R R R R R R R R R R R | Habitat: Ag, S, CF, AT, U, R, As | Ag, S, CF, AT, U, R, As | Ag, U, RPL, RpT, PJ, PP |
| | R R R R R b R R R R R R R R R | | | |
| | | Abundance: C | C | FC |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | Breeding | Migr. | Winter |
|--|--|--------------------------------------|--|--|
| | Habitat: | PJ | U | |
| Varied Thrush <u>Ixoreus naevius</u> | M W M — — — M W — M — — | | | |
| Acc. in 8, 14, 19, 23 | | | | |
| | | Abundance: R | R | |
| Gray Catbird <u>Dumetella carolinensis</u> | M B B B B B B B b B B B B B b b | B B b B b R B B R B B B R R | Habitat: RpL, RpT, MM, Sc | Breeding Migr. Winter RpL, RpT, MM, Sc |
| | | | | |
| | | Abundance: FC | FC | |
| Northern Mockingbird <u>Mimus polyglottos</u> | b b b R R N M B B B B B B B B B | B B B R b R R R B B R R | Habitat: Ag, S, PJ, RpL, RpT, CG | Breeding Migr. Winter Ag, S, PJ, RpL, CG |
| 9: One wintered, 1986-87 | | | | |
| | | Abundance: U-C | U | U |
| Sage Thrasher <u>Oreoscoptes montanus</u> | B B B M B B M B B B b B B B B | B M M B M M B M M M M M | Habitat: S | Breeding Migr. Winter S, PJ, RpT, U, CG |
| | | | | |
| | | Abundance: FC | FC | |
| Brown Thrasher <u>Toxostoma rufum</u> | M M R M R M M M M — M M | B B B B B R B B B M B B | Habitat: RpL, Ag | Breeding Migr. Winter RpL, Ag |
| Acc. in 8, 16 | | | | |
| | | R west | | |
| | | Abundance: FC east | R-FC | R |
| Bendire's Thrasher* <u>Toxostoma bendirei</u> | — — — — — — — B B — | — — — — B — | Habitat: undecided | Breeding Migr. Winter undecided |
| Acc. in 23, 27 | | | | |
| | | Und, 2 | | |
| | | Abundance: records | | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|--|--|------------|--------------------------------|----------------------------|-----------------------|--------------|
| <u>Curve-billed Thrasher</u> <u>Toxostoma curvirostre</u> | M M M M M R R N B R B | Habitat: | Breeding Sg, CG, PJ SgSd | Migr. CG, PJ | Winter CG, PJ | |
| Acc. in 10, 12, 14 18: One wintered, 1976 | | Abundance: | U | R | R | |
| <u>Water Pipit</u> <u>Anthus spinoletta</u> | M B B B W B B B M B B B B B B B | Habitat: | T | Breeding Aq, SgP, Ag | Migr. Ag | Winter Ag |
| Acc. in 1 | M M M b M M B M M | Abundance: | FC-C | C | FC | |
| <u>Sprague's Pipit#</u> <u>Anthus spragueii</u> | M M M M --- | Habitat: | | Breeding W/OG, CR | Migr. | Winter |
| Acc. in 4, 5, 12, 19 | | Abundance: | | R | | |
| <u>Bohemian Waxwing</u> <u>Bombycilla garrulus</u> | W W W W W W W W W W W W M W W W | Habitat: | | Breeding RpL | Migr. Ag, R, CF, U | Winter |
| Acc. in 14, 22 | W W W M W W W -- | Abundance: | Irr | Irr | | |
| <u>Cedar Waxwing</u> <u>Bombycilla cedrorum</u> | W b B R W N R R W B b N M R N W | Habitat: | R, Ag | Breeding R, Ag, PJ, U | Migr. U, R, Ag, PJ | Winter |
| Acc. in 22 | b N b W M b W N W N W W | Abundance: | U | FC | Irr | |
| <u>Phainopepla*</u> <u>Phainopepla nitens</u> | M M --- M | Habitat: | | Breeding RpT, Ag | Migr. | Winter |
| Acc. in 4, 5, 23 | --- | Abundance: | | R | | |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|--|------------------------------|---|----------------------------------|-------------------------------------|------------------------------------|
| <u>Northern Shrike</u> <u>Lanius excubitor</u> | <u>Acc. in 3</u> | <u>Habitat:</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| | | W W M W W W W W W W W W W W W W | W W W W W W W W W W W W | Ag | Ag, U, RpT, RpL, GL, PJ, S |
| | | | | Und | U-FC |
| <u>Loggerhead Shrike</u> <u>Lanius ludovicianus</u> | | <u>Habitat:</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| | | B B b B R B B R R B B R R B B | B B B B B B R B B B B B | SgP, MmP, CG, S, RpL, RpT, PJ | GL, S, Ag, U GL, S, Ag, U |
| | | | | FC-C | U-FC |
| <u>European Starling</u> <u>Sturnus vulgaris</u> | (Introduced) | <u>Habitat:</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| | | R R R R R R R R R R R R R R R R | R R R R R R R R R R R R | Ag, U, RpL, RpT | Ag, U, RpL, RpT |
| | | | | Ab | Ab |
| <u>White-eyed Vireo*</u> # <u>Vireo griseus</u> | <u>Acc. in 6, 11, 27, 28</u> | <u>Habitat:</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| | | — — M — — — — M — — — — — — M M | M | RpL | |
| | | | | R | |
| <u>Bell's Vireo</u> <u>Vireo bellii</u> | <u>Acc. in 4, 6, 19</u> | <u>Habitat:</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| | | — — M — M B — — — M — b — — — — M M | RpL, riparian thickets | RpL | |
| | | | | R-FC | R |
| <u>Gray Vireo</u> <u>Vireo vicinior</u> | | <u>Habitat:</u> | <u>Breeding</u> | <u>Migr.</u> | <u>Winter</u> |
| | | M B — B — — b b b — — b B b — — | PJ | PJ | |
| | | | | U | U |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|---|-------------------------------------|----------------------------|----------------|--|
| <u>Solitary Vireo</u> | B B B B | M M M | Habitat: | Breeding PP,As,R, Winter R,Ag,U PJ |
| <u>Vireo solitarius</u> | <u>B</u> <u>b</u> <u>B</u> <u>B</u> | <u>B</u> <u>M</u> <u>M</u> | | |
| Acc. in 13 | <u>b</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>M</u> <u>M</u> | | |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>M</u> <u>M</u> | | |
| | | | Abundance: FC | FC |
| <u>Yellow-throated Vireo#</u> | | | | |
| <u>Vireo flavifrons</u> | | | Habitat: | Breeding Migr. RpL,Ag |
| Acc. in 5, 11, 14, 19 | --- | M M | | |
| | --- | M | | |
| | --- | --- | Abundance: | R |
| <u>Warbling Vireo</u> | | | | |
| <u>Vireo gilvus</u> | B B b B | b b b | Habitat: | Breeding Migr. As,R U,R |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>b</u> <u>b</u> | | |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>b</u> <u>B</u> | | |
| | <u>B</u> <u>B</u> <u>B</u> <u>B</u> | <u>B</u> <u>b</u> <u>B</u> | | |
| | | | Abundance: FC | FC |
| <u>Philadelphia Vireo</u> | | | | |
| <u>Vireo philadelphicus</u> | --- | M M | Habitat: | Breeding Migr. RpL,RpT,U |
| Acc. in 5, 7, 14, 17, 19, 21, 23, 27 | --- | M M | | |
| | --- | M M | | |
| | --- | M M | | |
| | --- | M M | Abundance: | R |
| <u>Red-eyed Vireo</u> | | | | |
| <u>Vireo olivaceus</u> | M M B | b M M | Habitat: | Breeding Migr. RpL,U RpL,RpT, Ag,U |
| | <u>M</u> <u>M</u> <u>B</u> | <u>B</u> <u>M</u> <u>b</u> | | |
| | <u>M</u> <u>M</u> <u>b</u> | <u>B</u> <u>b</u> <u></u> | | |
| | <u>M</u> <u>M</u> <u>b</u> | <u>b</u> <u>M</u> <u></u> | | |
| Acc. in 1, 2, 10 | | | Abundance: R-U | R-U |
| <u>Blue-winged Warbler#</u> | | | | |
| <u>Vermivora pinus</u> | --- | M | Habitat: | Breeding Migr. RpL,Ag |
| Acc. in 11, 12, 13, 14, 19,28 | --- | M M M | | |
| | --- | M | | |
| | --- | --- | | |
| | --- | M | Abundance: | R |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

| | | | |
|----|-----------------|-----|----------------|
| Ab | - Abundant | R | - Rare |
| C | - Common | Und | - Undetermined |
| FC | - Fairly Common | Irr | - Irregular |
| U | - Unusual | | |

| | Breeding | Migr. | Winter |
|---|--|------------|--------------------------------|
| <u>Golden-winged Warbler</u> <u>Vermivora chrysoptera</u> | M M M M M M M M M M M M | Habitat: | RpL, Ag |
| Acc. in 6, 7, 12, 14, 19, 26, 27, 28 | | | |
| | | Abundance: | R |
| <u>Tennessee Warbler</u> <u>Vermivora peregrina</u> | M | Habitat: | SF, R, As, U, Ag |
| Acc. in 8, 9, 13, 15, 20, 21, 27 4: 2 males on territory, 7/13-25/81 11: 1 male on territory, 6/21-28/81 | | | |
| | | Abundance: | U |
| <u>Orange-crowned Warbler</u> <u>Vermivora celata</u> | M B B M M M M M M M M M B B B B M M M M M M M M B B B B M M M M M M M M M b b b b M M M M M M M M | Habitat: | Sc, PJ, R, As |
| Acc. in 1, 13 | | | Sc, PJ, R, As, PP, Ag |
| | | Abundance: | U |
| <u>Nashville Warbler</u> <u>Vermivora ruficapilla</u> | M | Habitat: | RpL, RpT, PJ, U |
| Acc. in 1, 2, 3, 8, 10, 15, 26 | | | |
| | | Abundance: | U |
| <u>Virginia's Warbler</u> <u>Vermivora virginiae</u> | b B b B M M M M M M M M B B B B B M M M M M M M M b B B B B M M M M M M M M B B b B B B M M M M M M M | Habitat: | Sc, RpT, MmP, As, DF, PJ |
| Acc. in 7 | | | RpT, RpL, MmP, Sc |
| | | Abundance: | FC-C |
| <u>Northern Parula</u> <u>Parula americana</u> | M | Habitat: | RpL |
| Acc. in 17, 21, 23, 24 | | | |
| 11: Unmated male on territory, 1981; Unmated female built nest, 1982 | | | |
| | | Abundance: | R |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | |
|---|--|------------|---|
| <u>Yellow Warbler</u> <u>Dendroica petechia</u> | B B B B B B B B B B B B B B b B B B B B B B B B B B B b | Habitat: | Breeding Migr. Winter RpL, RpT, R, U, Ag, S U, Ag |
| | | Abundance: | C FC-C |
| <u>Chestnut-sided Warbler</u> <u>Dendroica pensylvanica</u> | — — b M M M — — M B M — M — — M M M B — — — — M — — — — M | Habitat: | Breeding Migr. Winter RpL, Sc RpL, RpT, Ag, U |
| Acc. in 10, 16, 17, 18, 23 | | Abundance: | R U |
| <u>Magnolia Warbler</u> <u>Dendroica magnolia</u> | — — M M M M — — M M M M M M — — M — — M — M — — — — — — — M | Habitat: | Breeding Migr. Winter RpL, RpT |
| Acc. in 5, 9, 10, 13, 16, 28 | | Abundance: | R R |
| <u>Cape May Warbler</u> <u>Dendroica tigrina</u> | — — M M M — — W — — M — — — — — — — M | Habitat: | Breeding Migr. Winter RpL, U |
| Acc. in 6, 7, 14, 28 | | Abundance: | R |
| 11: One wintered, 1985-86 | | | |
| <u>Black-throated Blue Warbler</u> <u>Dendroica caerulescens</u> | — — M M M M — — M M — M — — M M — M — — — — — — — M | Habitat: | Breeding Migr. Winter Ag, RpL |
| Acc. in 7, 18, 21, 27 | | Abundance: | R |
| <u>Yellow-rumped Warbler</u> <u>Dendroica coronata</u> | B B B B M M M B b B B b M M b B B B B M M B B B B B M M | Habitat: | Breeding Migr. Winter CF, RpT CF, R, MmP, RpL, RpT U, As, Ag |
| | | Abundance: | C Ab R |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|-------------------------------------|---|--------------|-------------|---------------|--------|
| <u>Black-throated Gray Warbler</u> | B B M M M | Habitat: | Breeding PJ | Migr. PJ, Sc, | Winter |
| <u>Dendroica nigrescens</u> | <u>B</u> <u>B</u> <u>B</u> <u>M</u> <u>M</u> <u>M</u> | | | RpL, RPT | |
| Acc. in 6 | <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>M</u> | | | | |
| | <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>M</u> | | | | |
| | | Abundance: C | | U | |
| <u>Townsend's Warbler</u> | M M M M M M M | Habitat: | Breeding | Migr. | Winter |
| <u>Dendroica townsendi</u> | <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | CF, As, RpL, | |
| Acc. in 1, 3, 9, 15, 21 | <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | RpT, S | |
| | <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | | |
| | | Abundance: U | spring | FC fall | |
| <u>Hermit Warbler</u> | M | Habitat: | Breeding | Migr. | Winter |
| <u>Dendroica occidentalis</u> | <u>—</u> <u>—</u> <u>M</u> <u>—</u> <u>—</u> | | | RpL, RPT | |
| Acc. in 4, 11, 19 | <u>—</u> <u>—</u> <u>—</u> <u>M</u> <u>—</u> <u>—</u> | | | | |
| | <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> | | | | |
| | | Abundance: R | | | |
| <u>Black-throated Green Warbler</u> | M M M M M | Habitat: | Breeding | Migr. | Winter |
| <u>Dendroica virens</u> | <u>—</u> <u>M</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> | | | RpL | |
| Acc. in 6, 7, 9, 16, 27 | <u>—</u> <u>M</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> | | | | |
| | <u>—</u> <u>M</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> | | | | |
| | <u>—</u> <u>M</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> | | | | |
| | | Abundance: R | | | |
| <u>Blackburnian Warbler</u> | M M M M M | Habitat: | Breeding | Migr. | Winter |
| <u>Dendroica fusca</u> | <u>—</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | RpL, RpT, U | |
| Acc. in 5, 6, 7, 12, 21, 28 | <u>—</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | | |
| | <u>—</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | | |
| | <u>—</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | | |
| | | Abundance: R | | | |
| <u>Yellow-throated Warbler</u> | M M M | Habitat: | Breeding | Migr. | Winter |
| <u>Dendroica dominica</u> | <u>—</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | RpL | |
| Acc. in 4, 5, 11, 12, 14, 19 | <u>—</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | | |
| | <u>—</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | | |
| | <u>—</u> <u>—</u> <u>M</u> <u>M</u> <u>M</u> <u>M</u> | | | | |
| | | Abundance: R | | | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|--|---------------------------|----------|---------------------|---------|--------|
| <u>Grace's Warbler</u> | <u>Dendroica graciae</u> | Habitat: | Breeding | Migr. | Winter |
| | | | PP,Sc,PJ | RpL,RpT | |
| Acc. in 4, 8, | | | | | |
| | M b B B b | — — — | | | |
| | | | Abundance: FC | R-FC | |
| <u>Pine Warbler</u> | <u>Dendroica pinus</u> | Habitat: | Breeding | Migr. | Winter |
| | | | RpL,PP,Ag | | |
| Acc. in 5, 12, 14, 16, 19 | | | | | |
| | — — M M M — M | — — — | | | |
| | | | Abundance: R | | |
| <u>Prairie Warbler*</u> | <u>Dendroica discolor</u> | Habitat: | Breeding | Migr. | Winter |
| | | | RpL | | |
| Acc. in 4, 5, 11, 14 | | | | | |
| | — — M M — M | — — — | | | |
| | | | Abundance: R | | |
| <u>Palm Warbler</u> | <u>Dendroica palmarum</u> | Habitat: | Breeding | Migr. | Winter |
| | | | RpL,RpT,U | | |
| Acc. in 1, 5, 7, 10, 14, 16, 18, 20, 23, 24, 27 | | | | | |
| | M M M M M M M | — — — | | | |
| | | | Abundance: R | | |
| <u>Bay-breasted Warbler</u> | <u>Dendroica castanea</u> | Habitat: | Breeding | Migr. | Winter |
| | | | RpT,SF, 1 record | RpL,Ag | |
| Acc. in 5, 6, 7, 10, 25 | | | | | |
| | — — b M M M M | — — — | | | |
| | | | Abundance: Und | R | |
| <u>Blackpoll Warbler</u> | <u>Dendroica striata</u> | Habitat: | Breeding | Migr. | Winter |
| | | | RpL,RpT, Ag,U | | |
| Acc. in 13, 16, 21, 23, 24 | | | | | |
| | — — M M M M M M | — — — | | | |
| | | | Abundance: U | | |
| <hr/> | | | | | |
| STATUS | | | | | |
| R-Resident (breeds) | | | | | |
| N-Nonbreeder (resident) | | | | | |
| B-Definite breeder | | | | | |
| b-Likely breeder | | | | | |
| n-Non-breeder (summer) | | | | | |
| W-Winter visitor | | | | | |
| M-Migrant | | | | | |
| <hr/> | | | | | |
| ABUNDANCE | | | | | |
| Ab - Abundant | | | | | |
| C - Common | | | | | |
| FC - Fairly Common | | | | | |
| U - Unusual | | | | | |
| R - Rare | | | | | |
| Und - Undetermined | | | | | |
| Irr - Irregular | | | | | |

| | | | | |
|------------------------------------|---------------------|------------|--------------------|-------------------|
| <u>Black-and-white Warbler</u> | | Breeding | Migr. | Winter |
| <u>Mniotilla varia</u> | M M M M M M M M | Habitat: | RpL, Rpt, Sc, U | |
| Acc. in 5, 10, 13, 24, 26 | — — — — — — — — | | | |
| | | Abundance: | U | |
| <u>American Redstart</u> | | Breeding | Migr. | Winter |
| <u>Setophaga ruticilla</u> | M B B b M M B M M | Habitat: | R | R |
| Acc. in 13 | — — — — — — — — | | | |
| | | Abundance: | U-FC | FC-C |
| <u>Prothonotary Warbler</u> | | Breeding | Migr. | Winter |
| <u>Protonotaria citrea</u> | — — — M M M M M M | Habitat: | RpL | |
| Acc. in 6, 7, 11, 12, 14 19, 21 | — — — — — — — — | | | |
| | | Abundance: | R | |
| <u>Worm-eating Warbler</u> | | Breeding | Migr. | Winter |
| <u>Helmitheros vermivorus</u> | M M M M M M M M | Habitat: | RpL, Ag | |
| Acc. in 1, 5, 6, 14, 18, 27, 28 | — — — — — — — — | | | |
| | | Abundance: | R | |
| <u>Swainson's Warbler*</u> | | Breeding | Migr. | Winter |
| <u>Limnothlypis swainsonii</u> | — — — M — — — — | Habitat: | RpL | |
| Acc. in 4, 19 | — — — — — — — — | | | |
| | | Abundance: | R | |
| <u>Ovenbird</u> | | Breeding | Migr. | Winter |
| <u>Seiurus aurocapillus</u> | — — — B B M M B M M | Habitat: | PP, RpL | Sc, RpL, Ag, U |
| Acc. in 13, 21 | — — — B B M M B M M | | | |
| | | Abundance: | U | U-FC |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | Breeding | Migr. | Winter |
|----------------------------------|--------------------------------------|------------|----------------------------------|
| <u>Northern Waterthrush</u> | | | |
| <u>Seiurus noveboracensis</u> | M M M M M M M | Habitat: | RpL, RpT, Ag, U, Ms |
| Acc. in 1, 3, 8, 9, 10, 26 | M M M M M M M M M M M M M M | | |
| | | Abundance: | U |
| <u>Louisiana Waterthrush*#</u> | | | |
| <u>Seiurus motacilla</u> | ----- M | Habitat: | RpL |
| Acc. in 14, 27 | ----- M ----- M | | |
| | | Abundance: | R |
| <u>Kentucky Warbler*</u> | | | |
| <u>Oporornis formosus</u> | --- M M M M | Habitat: | RpL, U |
| Acc. in 7, 10, 11, 21, 23, 26 | --- M M M M --- M M M M | | |
| | | Abundance: | R |
| <u>Connecticut Warbler*</u> | | | |
| <u>Oporornis agilis</u> | ----- M | Habitat: | RpL |
| Acc. in 5 | ----- M | | |
| | | Abundance: | Und |
| <u>Mourning Warbler*</u> | | | |
| <u>Oporornis philadelphia</u> | ----- M M ----- | Habitat: | 1 record only |
| Acc. in 4, 12, 19 | ----- M M ----- | | |
| | | Abundance: | Und |
| <u>MacGillivray's Warbler</u> | | | |
| <u>Oporornis tolmiei</u> | B B b B b M M | Habitat: | R, Sc, As R, Sc, U, PJ, Ag |
| Acc. in 7, 21 | B B B B B M M B B B B B M M | | |
| | | Abundance: | FC |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

Common Yellowthroat

Geothlypis trichas

| B | B | b | B | B | B | B | B | B | B | Habitat: | Breeding | Migr. | Winter |
|---|---|---|---|---|---|---|---|---|---|----------|----------|-------|--------|
| — | — | — | — | — | — | — | — | — | — | R,Ms | R,Ms,Ag | | |
| — | — | — | — | — | — | — | — | — | — | | | | |
| — | — | — | — | — | — | — | — | — | — | | | | |
| — | — | — | — | — | — | — | — | — | — | | | | |

Abundance: C

C

Hooded Warbler

*Wilsonia citrina*Acc. in 5, 7, 14, 20, 21,
25, 27

| — | — | M | M | M | M | Habitat: | Breeding | Migr. | Winter |
|---|---|---|---|---|---|----------|----------|-------|--------|
| — | — | — | — | — | — | RpL,U | | | |
| — | — | — | — | — | — | | | | |
| — | — | — | — | — | — | | | | |
| — | — | — | — | — | — | | | | |

Abundance: R

Wilson's Warbler

Wilsonia pusilla

| B | B | B | B | M | M | M | Habitat: | Breeding | Migr. | Winter |
|---|---|---|---|---------|-----|----------|----------|----------|-------|--------|
| — | — | — | — | AT,RpH, | RpT | R,S,Ag,U | | | | |
| — | — | — | — | | | | | | | |
| — | — | — | — | | | | | | | |
| — | — | — | — | | | | | | | |

Abundance: FC-Ab

C

Canada Warbler

*Wilsonia canadensis*Acc. in 4, 5, 11, 12, 19
11: One territorial singing
male 6/24-7/11/84Otherwise, no records since
1976

| — | — | M | M | — | — | Habitat: | Breeding | Migr. | Winter |
|---|---|---|---|----------|---|----------|----------|-------|--------|
| — | — | — | — | RpL,Ag,U | | | | | |
| — | — | — | — | | | | | | |
| — | — | — | — | | | | | | |
| — | — | — | — | | | | | | |

Abundance: R

Painted Redstart*

Myioborus pictus

Acc. in 4, 11, 16

| — | — | M | — | — | — | Habitat: | Breeding | Migr. | Winter |
|---|---|---|---------|---|---|----------|----------|-------|--------|
| — | — | — | RpL,RpT | | | | | | |
| — | — | — | | | | | | | |
| — | — | — | | | | | | | |
| — | — | — | | | | | | | |

Abundance: R

Yellow-breasted Chat

Icteria virens

Acc. in 10

| B | b | b | B | b | b | B | b | B | b | Habitat: | Breeding | Migr. | Winter |
|---|---|---|---|---|---|---|---|---|---|----------|----------|-------|--------|
| — | — | — | — | — | — | — | — | — | — | R,S | R,S,Ag,U | | |
| — | — | — | — | — | — | — | — | — | — | | | | |
| — | — | — | — | — | — | — | — | — | — | | | | |
| — | — | — | — | — | — | — | — | — | — | | | | |

Abundance: FC

U

STATUS

- R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

- | | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|--|--|---------------------|---------|------------|
| <u>Hepatic Tanager*</u> # | | Breeding | Migr. | Winter |
| <u>Piranga flava</u> | | PP | PP, RpT | |
| Acc. in 8, 11, 14 | | | | |
| | M --- M --- M ----- B ----- | | | |
| | | Abundance: Und | R | |
| <u>Summer Tanager</u> | | Breeding | Migr. | Winter |
| <u>Piranga rubra</u> | | RpL, RpT, Ag, U | | |
| Acc. in 5, 6, 8, 10, 13, 22, 25, 27, 28 | | | | |
| | M --- M M M M M M M M --- M M M M | Habitat: | | |
| | | | | |
| | | Abundance: | R | |
| <u>Scarlet Tanager</u> | | Breeding | Migr. | Winter |
| <u>Piranga olivacea</u> | | RpL, RpT | | |
| Acc. in 5, 7, 9, 12, 14, 16, 18, 19, 21, 26 | | | | |
| | M --- M M M M M M M M M M M M M M | Habitat: | | |
| | | | | |
| | | Abundance: | R | |
| <u>Western Tanager</u> | | Breeding | Migr. | Winter |
| <u>Piranga ludoviciana</u> | | PP, Sc, As, SF | R, U | |
| 5-One summered, 1985 | | | | |
| | b B b B n M M B B B B B M M b B B B B M M B B B B B M M | Habitat: | | |
| | | | | |
| | | Abundance: | FC | |
| <u>Northern Cardinal</u> | | Breeding | Migr. | Winter |
| <u>Cardinalis cardinalis</u> | | RpL | RpL, U | RpL, Ag, U |
| Acc. in 21, 25, 28 | | | | |
| | W W M M b R W W W W M M M M M | Habitat: | | |
| | | | | |
| 18: One wintered, 1985-86 | | Abundance: | R | R |
| <u>Rose-breasted Grosbeak</u> | | Breeding | Migr. | Winter |
| <u>Pheucticus ludovicianus</u> | | RpL, RpT, U, Ag, | | |
| Acc. 2, 5, 13 | | | | |
| | M | Habitat: | | |
| | | | | |
| | | Abundance: | U | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|--|--|----------------------------------|------------|-------------------------------|----------------------|--------|
| <u>Black-headed Grosbeak</u> <u>Pheucticus melanocephalus</u> | B B M B B B B B B B B B B B B B | b M B B b b B B M B B M | Habitat: | Breeding PP, Sc, PJ, As, R | Migr. R, U | Winter |
| | | | Abundance: | FC | FC | |
| <u>Blue Grosbeak</u> <u>Guiraca caerulea</u> | B M - B B - B B B M B B B M b | b B B B B B B B B B B B | Habitat: | Breeding RpL, RpT, PJ, CG | Migr. RpL, RpT, S | Winter |
| | | | Abundance: | U north | FC south | FC |
| <u>Lazuli Bunting</u> <u>Passerina amoena</u> | b B b B B B B B B B B B b B b B | B M M B M M B M M B B M | Habitat: | Breeding RpL, RpT, Sc, Sa, PJ | Migr. RpL, RpT, S, U | Winter |
| | | | Abundance: | FC-C | FC | |
| <u>Indigo Bunting</u> <u>Passerina cyanea</u> | M - B M B B B M M B b B - B | b b b B B B B B B B B B | Habitat: | Breeding RpL, RpT, S | Migr. RpL, RpT, U | Winter |
| Acc. in 1, 9, 16 | | | Abundance: | U | U | |
| <u>Painted Bunting</u> <u>Passerina ciris</u> | - - - M - - - - - - M - M - - - - M - - - - - | | Habitat: | Breeding | Migr. | Winter |
| Acc. in 4, 19, 21, 28 | | | | RpL | | |
| | | | Abundance: | | R | |
| <u>Dickcissel</u> <u>Spiza americana</u> | b - M M - B M M - b - - - - | b B B B B B b B B B B | Habitat: | Breeding GL, Cr | Migr. GL | Winter |
| Acc. in 8, 15, 16 | | | Abundance: | Irr | R | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|--------------------------------|----------------|--------------|------------|--------------|---------------|-----------|
| <u>Green-tailed Towhee</u> | B B B B | M M M | Habitat: | Breeding | Migr. | Winter |
| <u>Pipilo chlorurus</u> | <u>B B B B</u> | <u>B M M</u> | | RpL, RpT, S, | R, S, MmP, | |
| Acc. in 13 | <u>B B B B</u> | <u>B M</u> | | MmP, PJ | CG | |
| | <u>B B B B</u> | <u>B b M</u> | | | | |
| | | | Abundance: | FC | FC | |
| <u>Rufous-sided Towhee</u> | | | | | | |
| <u>Pipilo erythrophthalmus</u> | B B b R | M M b | Habitat: | Breeding | Migr. | Winter |
| | <u>R R B R</u> | <u>B M W</u> | | RpL, RpT, S | R, S, U, CG | S, U |
| | <u>R R b R</u> | <u>R R W</u> | | | | |
| | <u>R R B R</u> | <u>R B R</u> | | | | |
| | | | Abundance: | FC | FC | U |
| <u>Brown Towhee</u> | | | | | | |
| <u>Pipilo fuscus</u> | M — — M | M — b — | Habitat: | Breeding | Migr. | Winter |
| | <u>M — — M</u> | <u>R R M</u> | | PJ, S, CG | S, CG | PJ, S, CG |
| Acc. in 4, 5, 8, 11, 17 | <u>M — — R</u> | <u>R R R</u> | | | | |
| | | | Abundance: | FC | R-FC | FC |
| <u>Cassin's Sparrow</u> | | | | | | |
| <u>Aimophila cassinii</u> | — — — M | B B b | Habitat: | Breeding | Migr. | Winter |
| | <u>b B B</u> | <u>B B B</u> | | SgP, S, SS | SgP, S, SS | |
| Acc. in 11 | <u>B B B</u> | <u>B B B</u> | | | | |
| | | | Abundance: | FC-C | FC | |
| <u>Rufous-crowned Sparrow</u> | | | | | | |
| <u>Aimophila ruficeps</u> | M — — M | — — — | Habitat: | Breeding | Migr. | Winter |
| | <u>M — — M</u> | <u>M M</u> | | Sc | Sc | |
| Acc. in 8, 11, 19 | <u>b B B</u> | <u>B B B</u> | | | | |
| | | | Abundance: | FC | R | |
| <u>American Tree Sparrow</u> | | | | | | |
| <u>Spizella arborea</u> | W W W W | W W W | Habitat: | Breeding | Migr. | Winter |
| | <u>W W W W</u> | <u>W W W</u> | | R | RpL, RpT, | |
| | <u>W W W W</u> | <u>W W W</u> | | | GL, Ag, Sa, U | |
| | <u>W W W W</u> | <u>W W W</u> | | | | |
| | | | Abundance: | FC | FC | FC-Ab |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|--|--|------------|------------------------------|---------------------------|--------|
| <u>Chipping Sparrow</u> <u>Spizella passerina</u> | B B b B B B M | Habitat: | Breeding R,PP,PJ,S | Migr. R,GL,U, Sa,Sc,Ag | Winter |
| 23: One wintered, Jan. 1982 | B B B B B b M B B B B B b M B R B B B B M | | | | |
| | | Abundance: | FC-C | FC-Ab | |
| <u>Clay-colored Sparrow</u> <u>Spizella pallida</u> | M M M M M M M M | Habitat: | Breeding | Migr. | Winter |
| Acc. in 15 | M M M M M M M M M M M M M M M M — — — — M M M M | | | R,GL,Ag, U | |
| | | Abundance: | | FC | |
| <u>Brewer's Sparrow</u> <u>Spizella breweri</u> | B B B B B B b | Habitat: | Breeding S,SgSD, SS | Migr. GL,R,S,PJ | Winter |
| 15: Two wintered, 1984-85 | B B B B B b B R B B b B b M B B B B b M M | | | | |
| | | Abundance: | C | FC | |
| <u>Field Sparrow</u> <u>Spizella pusilla</u> | W W M M b | Habitat: | Breeding RpL | Migr. RpL | Winter |
| Acc. in 5, 6, 7, 8, 9, 15 | M — M M M M b M — — — M M b — — — — b | | | | |
| 4: One wintered, 1983-84 | | Abundance: | R | R | |
| 9: One wintered, 1986-87 | | | | | |
| <u>Vesper Sparrow</u> <u>Pooecetes gramineus</u> | B B B B B B B | Habitat: | Breeding GL,S,PJ | Migr. GL,S | Winter |
| | B B B B B B b B B B B B b b B B B B B B B | | | | |
| | | Abundance: | U-C | FC-C | |
| <u>Lark Sparrow</u> <u>Chondestes grammacus</u> | B b B B B B B | Habitat: | Breeding GL,S,RpL, RpT,PJ | Migr. GL,As,S | Winter |
| | B M B B B B B B B b B B B B B B B B B B B | | | | |
| | | Abundance: | FC-Ab | FC-Ab | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|-------------------------------|----------------------------------|--------------------------------------|------------|----------------------|----------------------|--------|
| <u>Black-throated Sparrow</u> | <u>Amphispiza bilineata</u> | M M M M — — | Habitat: | Breeding | Migr. | Winter |
| | | <u>B b M M M M — —</u> | | PJ, S, SgSD, CG | Sa, SgSD, CG | |
| Acc. in 4, 5, 10 | | <u>B M M B B — —</u> | | | | |
| | | <u>B B — — — b M</u> | | | | |
| | | | Abundance: | U-C | R-U | |
| <u>Sage Sparrow</u> | <u>Amphispiza belli</u> | B B B B M M — | Habitat: | Breeding | Migr. | Winter |
| | | <u>B B B — M — —</u> | | Sa, SgSD | S, SgSD | |
| Acc. in 5, 6, 18, 20, 24 | | <u>B B B M M M —</u> | | | | |
| | | <u>B B M B — — —</u> | | | | |
| | | | Abundance: | U-FC | R | |
| <u>Lark Bunting</u> | <u>Calamospiza melanocorys</u> | b b B B B B B B | Habitat: | Breeding | Migr. | Winter |
| | | <u>b b M B b B B B</u> | | GL, Sa | GL, Ag, SaRb | |
| | | <u>M M b b B B B B</u> | | | | |
| | | <u>M M M b B B B B</u> | | | | |
| | | | Abundance: | R-FC west Ab east | R-FC west Ab east | |
| <u>Savannah Sparrow</u> | <u>Passerculus sandwichensis</u> | b B B B B B b | Habitat: | Breeding | Migr. | Winter |
| | | <u>b b B B B — M</u> | | GL, Ms | GL | |
| | | <u>b B B B b b M</u> | | | | |
| | | <u>M B B B b M B</u> | | | | |
| | | | Abundance: | FC | FC | |
| <u>Baird's Sparrow</u> | <u>Ammodramus bairdii</u> | M — — M M M M | Habitat: | Breeding | Migr. | Winter |
| | | <u>M — — M M M M</u> | | | GL | |
| Acc. in 16, 28 | | <u>M — — M M M M</u> | | | | |
| | | <u>M — — — M M M</u> | | | | |
| | | | Abundance: | — | R | |
| <u>Grasshopper Sparrow</u> | <u>Ammodramus savannarum</u> | b M b B B B B B | Habitat: | Breeding | Migr. | Winter |
| | | <u>b M b B B B B B</u> | | GL, TgP, MxP | GL | |
| Acc. in 16 | | <u>M — — B B B B B</u> | | | | |
| | | <u>M — — b B B B B</u> | | | | |
| | | | Abundance: | U-Irr | R | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | | |
|------------------------------|------------------------------|--|------------|--|-------|--------|
| <u>Henslow's Sparrow</u> | <u>Ammodramus henslowii</u> | M | Habitat: | Breeding | Migr. | Winter |
| Acc. in 5 | | ----- | | | | |
| | | ----- | | | | |
| | | ----- | | | | |
| | | ----- | | | | |
| | | | Abundance: | | | |
| <u>Le Conte's Sparrow*</u> | <u>Ammodramus lecontei</u> | M | Habitat: | Breeding | Migr. | Winter |
| Acc. in 4 | | ----- | | | | |
| | | ----- | | | | |
| | | ----- | | | | |
| | | | Abundance: | R | | |
| <u>Sharp-tailed Sparrow*</u> | <u>Ammodramus caudacutus</u> | | Habitat: | Breeding | Migr. | Winter |
| Acc. in 17, 19 | | ----- | | SgP, 2 records only | | |
| | | ----- | | | | |
| | | ----- | | | | |
| | | | Abundance: | R | | |
| <u>Fox Sparrow</u> | <u>Passerella iliaca</u> | b b R M B B N B M M M b b B W B b M | Habitat: | Breeding RpT, RpH Migr. R Winter RpL, RpT | | |
| Acc. in 13, 24 | | ----- | | | | |
| | | ----- | | | | |
| | | ----- | | | | |
| <u>Song Sparrow</u> | <u>Melospiza melodia</u> | B R R R R R M R R R R R R R R R R R R M R R R R R R R | Habitat: | Breeding R, Ms, U Migr. R, Ms, U Winter R, Ms, U | | |
| Acc. in 21 | | ----- | | | | |
| | | ----- | | | | |
| | | ----- | | | | |
| | | | Abundance: | C | FC | |
| <u>Lincoln's Sparrow</u> | <u>Melospiza lincolnnii</u> | M M B B M M M B B B B M M M b B B B b M M B b B B B M M | Habitat: | Breeding AT, MmP, Rph, RpT Migr. R, U, Ms | | |
| Acc. in 13 | | ----- | | | | |
| | | ----- | | | | |
| | | ----- | | | | |
| | | | Abundance: | FC | | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|-------------------------------|--|------------|----------|------------------------|-------------------|
| <u>Swamp Sparrow</u> | <u>Melospiza georgiana</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 8, 15, 16, 28 | | | | Ms, RpL | Ms, RpL |
| | M — — M W W M M — — W W — M M — W W — — — M — M M | | | | |
| | | Abundance: | R | U | |
| <u>White-throated Sparrow</u> | <u>Zonotrichia albicollis</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 9, 13, 15, 21 | | | | RpL, RpT, U | RpL, U |
| | M W W W M M M W M W W W M M M M M W W W M M M — — — M | | | | |
| | | Abundance: | R | U | |
| <u>Golden-crowned Sparrow</u> | <u>Zonotrichia atricapilla</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. 4, 7, 23 | | | | Sc, RpT | |
| | — — W — M — — — — — W — — — | | | | |
| | | Abundance: | R | | |
| <u>White-crowned Sparrow</u> | <u>Zonotrichia leucophrys</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 15 | | | AT, RpH | R, S, U, PP, PJ, Ag | R, U, Ag |
| | B R R R W W W R B B R W W W R R B R R W W R R B R R W W | | | | |
| | | Abundance: | FC-C | C | C |
| <u>Harris' Sparrow</u> | <u>Zonotrichia querula</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 15 | | | | RpL, RpT, U, S | RpL, RpT, U, S |
| | W W W W W W W W W W W W W W M M W W W W W W W M W W W W | | | | |
| | | Abundance: | U | U | U west FC east |
| <u>Dark-eyed Junco</u> | <u>Junco hyemalis</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 15 | | | CF, As | R, U, As, CF | U, R, PP, S |
| | R B R R W W W R R R R R W W R R R R R W W R R R R R W W | | | | |
| | | Abundance: | FC-C | FC-C | FC-Ab |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|-----------------------------------|------------------------------|-------------------|--|--|---------------|
| <u>McCown's Longspur</u> | <u>Calcarius mccownii</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 7, 8, 20 | | | SgP | SgP, MmP | |
| | | | <u>M</u> — <u>M</u> <u>B</u> <u>B</u> <u>B</u> <u>M</u> | <u>M</u> — <u>M</u> <u>B</u> <u>M</u> <u>M</u> — | |
| | | | — — — <u>M</u> <u>M</u> <u>M</u> — | — — — <u>M</u> <u>M</u> — | |
| | | <u>Abundance:</u> | FC | U | |
| <u>Lapland Longspur</u> | <u>Calcarius lapponicus</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 2 | | | GL, Cr | GL, Cr | |
| | | | <u>W</u> <u>W</u> <u>W</u> <u>W</u> <u>W</u> <u>W</u> | <u>W</u> <u>W</u> <u>W</u> <u>W</u> <u>W</u> — | |
| | | | <u>W</u> <u>M</u> <u>W</u> <u>W</u> <u>W</u> <u>W</u> — | <u>W</u> <u>W</u> — <u>W</u> <u>W</u> | |
| | | <u>Abundance:</u> | U-C | U-Ab | |
| <u>Chestnut-collared Longspur</u> | <u>Calcarius ornatus</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 14, 17 | | | SgP, MXP | SgP, MmP | |
| | | | <u>M</u> <u>M</u> <u>M</u> — <u>B</u> <u>B</u> <u>M</u> | <u>M</u> — <u>M</u> — | |
| | | | — — — <u>M</u> <u>M</u> — <u>M</u> — | — — — <u>M</u> <u>M</u> — | |
| | | <u>Abundance:</u> | FC | U-FC | |
| <u>Snow Bunting</u> | <u>Plectrophenax nivalis</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 10, 15, 24 | | | Sa, GL | | |
| | | | <u>W</u> <u>W</u> <u>W</u> <u>W</u> <u>W</u> <u>W</u> | <u>W</u> <u>W</u> <u>W</u> <u>W</u> — — | |
| | | | <u>W</u> <u>W</u> <u>W</u> <u>W</u> <u>W</u> — — | <u>W</u> — — <u>W</u> — — | |
| | | <u>Abundance:</u> | R | | |
| <u>Bobolink</u> | <u>Dolichonyx oryzivorus</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| Acc. in 15, 20 | | | GL, Cr-hayfields, fallow fields | Gl, Cr | |
| | | | <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> | <u>B</u> <u>b</u> <u>B</u> <u>b</u> — — | |
| | | | <u>B</u> <u>b</u> <u>M</u> <u>B</u> <u>B</u> — — | <u>M</u> <u>b</u> <u>B</u> <u>b</u> <u>M</u> <u>M</u> — | |
| | | <u>Abundance:</u> | R | | |
| <u>Red-winged Blackbird</u> | <u>Agelaius phoeniceus</u> | <u>Habitat:</u> | Breeding | Migr. | Winter |
| | | | Ms, Ag, GL, Sa | Ms, Ag, R, Sa | Ms, Ag, U, Sa |
| | | | <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> | <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> | |
| | | | <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> | <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> | |
| | | <u>Abundance:</u> | Ab | Ab | C |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

Eastern Meadowlark*Sturnella magna

Acc. in 13, 28

7: No longer nesting here, as field was burned, eliminating habitat

| | b | Habitat: | Breeding | Migr. | Winter |
|-------|---|----------|--------------|-------|--------|
| ----- | M | | SgP, TgP, Cr | | |
| ----- | M | | | | |
| ----- | M | | | | |
| ----- | | | | | |

Abundance: U

Western MeadowlarkSturnella neglecta

| | | | Habitat: | Breeding | Migr. | Winter |
|---------|-------|--|----------|--------------|-------|--------|
| R R B R | R R R | | | Sg, MmP, Sa, | S, GL | |
| R B B R | R b R | | | SgSD, PJ, | | |
| R R R R | R R R | | | Ag, SaRb, | | |
| B R R R | R R R | | | Cr, CG, SS | | |

Abundance: C-Ab C FC

Yellow-headed BlackbirdXanthocephalus xanthocephalus

| | | | Habitat: | Breeding | Migr. | Winter |
|---------|-------|--|----------|----------|--------------|--------|
| B B B B | B B B | | | Ms | Ms, Ag, RpL, | |
| B B B B | B b B | | | U | | |
| B B B B | B B b | | | | | |
| B B B B | B b M | | | | | |

Abundance: C C-Ab

Rusty BlackbirdEuphagus carolinus

Acc. in 7, 16, 27

| | | | Habitat: | Breeding | Migr. | Winter |
|-------|-------|--|----------|----------|-------|-----------|
| --- W | W W M | | | | | RpL, RpT, |
| W | W W W | | | | | Ag |
| M W | W W W | | | | | |
| --- | M | | | | | |

Abundance: R-U

Brewer's BlackbirdEuphagus cyanocephalus

| | | | Habitat: | Breeding | Migr. | Winter |
|---------|-------|--|----------|-------------|------------|------------|
| B B B R | R B b | | | R, Ag, MmP, | R, Ag, Sa, | RpL, Ag, U |
| R B B B | R b W | | | U | Sc | |
| B B B B | R R N | | | | | |
| B B R B | B R B | | | | | |

Abundance: FC-C C-Ab R-FC

Great-tailed GrackleQuiscalus mexicanusAcc. in 5, 11, 12, 15, 16,
18, 20, 25
Expanding

26: One wintered, 1979-80

| | | |
|---------|-------|--|
| B - B M | M -- | |
| M M B b | B M M | |
| B B R b | W - B | |

Abundance: R

R

STATUS

R-Resident (breeds)

N-Nonbreeder (resident)

B-Definite breeder

b-Likely breeder

n-Non-breeder (summer)

W-Winter visitor

M-Migrant

ABUNDANCE

Ab - Abundant

R - Rare

C - Common

Und - Undetermined

FC - Fairly Common

Irr - Irregular

U - Unusual

| Common Grackle | B | B | B | R | B | B | B | Habitat: | Breeding | Migr. | Winter |
|----------------------------|-----------|-----------|----------|----------|----------|----------|----------|------------|------------|------------------|--------|
| <u>Quiscalus quiscula</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | | R,Ag,U | RpL,RpT, Ag,U | |
| Acc. in 16 | b | b | b | b | b | b | b | | | | |
| Expanding on Western Slope | b | b | M | M | b | b | b | | | | |
| 4: Two wintered, 1982-83 | b | b | M | M | b | b | b | | | | |
| | | | | | | | | Abundance: | C | C | |
| Brown-headed Cowbird | B | B | B | B | B | b | B | Habitat: | Breeding | Migr. | Winter |
| <u>Molothrus ater</u> | <u>B</u> | <u>B</u> | <u>b</u> | <u>B</u> | <u>B</u> | <u>b</u> | <u>B</u> | | Ag,CF,S,U, | R,Ag | |
| | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | | MmP,R,As, | | |
| | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | | Ms | | |
| | | | | | | | | Abundance: | FC | FC | |
| Orchard Oriole | B | B | B | B | B | B | B | Habitat: | Breeding | Migr. | Winter |
| <u>Icterus spurius</u> | <u>--</u> | <u>--</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | | RpL,Ag | RpL,RpT,U, | |
| Acc. in 17 | <u>M</u> | <u>M</u> | <u>B</u> | <u>b</u> | <u>B</u> | <u>b</u> | <u>B</u> | | Ag | | |
| | <u>M</u> | <u>M</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | | | | |
| | <u>b</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | | | | |
| | | | | | | | | Abundance: | FC east | U-FC | |
| Northern Oriole | B | B | B | B | B | B | B | Habitat: | Breeding | Migr. | Winter |
| <u>Icterus galbula</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | | RpL,RpT, | RpL,RpT, | |
| | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | | Ag,U | Ag,U | |
| | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | <u>B</u> | | | | |
| | | | | | | | | Abundance: | FC | FC | |
| Scott's Oriole | B | M | M | M | M | M | M | Habitat: | Breeding | Migr. | Winter |
| <u>Icterus parisorum</u> | <u>B</u> | <u>M</u> | <u>M</u> | <u>M</u> | <u>M</u> | <u>M</u> | <u>M</u> | | DS,Sa | R | |
| Acc. in 10, 11, 15 | <u>M</u> | <u>M</u> | <u>M</u> | <u>M</u> | <u>M</u> | <u>M</u> | <u>M</u> | | | | |
| | <u>b</u> | <u>b</u> | <u>b</u> | <u>b</u> | <u>b</u> | <u>b</u> | <u>b</u> | | | | |
| | | | | | | | | Abundance: | Und | R | |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | |
|------------------------------------|------------------------|------------------------|------------------------|------------------------|
| Brambling* | | Breeding | Migr. | Winter |
| <u>Fringilla montifrigilla</u> | M | | | Urb |
| Acc. in 4, 19 | | | | |
| | M | | | |
| | | | | |
| | | Abundance: | R | |
| Rosy Finch | | Breeding | Migr. | Winter |
| <u>Leucosticte arctoa</u> | W R R R W W M | T,cliffs | U,Sa,CF, dirt banks | U,Sa,CF, dirt banks |
| Acc. in 7, 14 | W R R R R W W | AT-cliffs, | AT-cliffs, | |
| | W R R R R R | MmP, SgP, SgSD, SgM | MmP, SgP, SgSD, SgM | |
| | R R R R b | | | |
| | | Abundance:FC | U-Ab | U-Ab |
| Pine Grosbeak | | Breeding | Migr. | Winter |
| <u>Pinicola enucleator</u> | W W B R SF | SF | CF | CF |
| Acc. in 21 | W W B R R | | | |
| | b b R R R | | | |
| | R R R R R | | | |
| | R R R R R | Abundance:FC | U | U-FC |
| Purple Finch | | Breeding | Migr. | Winter |
| <u>Carpodacus purpureus</u> | W M M M | | | RpL,U |
| Acc. in 6, 7, 14, 18 21, 23, 27 | W M W M | | | |
| | W M M | | | |
| 23: One wintered, 1974-75 | | Abundance: | U | |
| Cassin's Finch | | Breeding | Migr. | Winter |
| <u>Carpodacus cassinii</u> | R B R R M CF, Sc, | CF, U | | RpL, RpT, PP, U |
| Acc. in 6, 28 | R R R R W W RpT | | | |
| | b R R B R W | | | |
| | R R D R B M M | | | |
| | | Abundance:FC | FC | FC |
| House Finch | | Breeding | Migr. | Winter |
| <u>Carpodacus mexicanus</u> | B B R R R R N U, SaRb, | U, RpL, | U, RpL, RpT, | |
| | R B B R R b W PJ | RpT, Sa, | Sa, Ag, | |
| | R R R R R R SgSD, CG | SgSD, CG | SgSD, CG | |
| | R R R R B B R | | | |
| | | Urban-Ab | Urban-Ab | Urban-Ab |
| | Abundance:U-C | C | U-C | U-C |

STATUS

R-Resident (breeds)
 N-Nonbreeder (resident)
 B-Definite breeder
 b-Likely breeder
 n-Non-breeder (summer)
 W-Winter visitor
 M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | | | | |
|--|--|----------------------------|----------------------|--------------------------|------------------------|
| <u>Red Crossbill</u> <u>Loxia curvirostra</u> | b R R R M M M | Habitat: | Breeding CF | Migr. CF | Winter CF, RpL |
| Acc. in 6, 7, 27 | <u>R</u> <u>D</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> | | | | |
| | <u>B</u> <u>b</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> | <u>R</u> <u>W</u> <u>—</u> | | | |
| | <u>b</u> <u>R</u> <u>N</u> <u>R</u> <u>R</u> <u>R</u> | <u>b</u> <u>M</u> <u>M</u> | | | |
| | | | Abundance: Irr-Ab | Irr-FC | Irr-FC |
| <u>White-winged Crossbill</u> <u>Loxia leucoptera</u> | <u>N</u> <u>N</u> <u>M</u> <u>—</u> | Habitat: | Breeding CF | Migr. CF | Winter CF |
| Acc. in 6, 12, 23, 24, 26 | <u>W</u> <u>W</u> <u>N</u> <u>M</u> <u>—</u> | | | | |
| | <u>—</u> <u>—</u> <u>N</u> <u>N</u> <u>—</u> | | | | |
| | <u>—</u> <u>M</u> <u>M</u> <u>M</u> <u>—</u> | | | | |
| | | | Abundance: R | R | |
| <u>Common Redpoll</u> <u>Carduelis flammea</u> | <u>M</u> <u>W</u> <u>W</u> <u>W</u> <u>W</u> | Habitat: | Breeding | Migr. | Winter |
| Acc. in 1, 10, 26 | <u>W</u> <u>—</u> <u>M</u> <u>W</u> <u>W</u> | | | | GL, S, U |
| | <u>—</u> <u>W</u> <u>W</u> <u>W</u> <u>W</u> | | | | |
| | <u>—</u> <u>W</u> <u>—</u> <u>—</u> <u>M</u> | | | | |
| | | | Abundance: Irr | | |
| <u>Pine Siskin</u> <u>Carduelis pinus</u> | <u>N</u> <u>B</u> <u>B</u> <u>R</u> <u>n</u> <u>W</u> <u>W</u> | Habitat: | Breeding CF, R, Sc | Migr. CF, R, Sc, U, MmP | Winter RpL, U, Sc, PP |
| | <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>W</u> <u>W</u> | | | | |
| | <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>W</u> <u>N</u> | | | | |
| | <u>R</u> <u>R</u> <u>R</u> <u>b</u> <u>R</u> <u>W</u> <u>W</u> | | | | |
| | | | Abundance: FC | AB | C |
| <u>Lesser Goldfinch</u> <u>Carduelis psaltria</u> | <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>M</u> | Habitat: | Breeding RpL, RpT, S | Migr. RpL, RpT, S, CG, U | Winter RpL, U |
| | <u>R</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>—</u> | | | | |
| | <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>—</u> | | | | |
| | <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> | | | | |
| | | | Abundance: FC | FC | R |
| <u>American Goldfinch</u> <u>Carduelis tristis</u> | <u>B</u> <u>R</u> <u>B</u> <u>R</u> <u>R</u> <u>R</u> <u>b</u> | Habitat: | Breeding RpL, RpT | Migr. RpL, RpT, Ag, U | Winter RpL, RpT, Ag, U |
| | <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> | | | | |
| | <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> <u>R</u> | | | | |
| | <u>R</u> <u>R</u> <u>b</u> <u>b</u> <u>b</u> <u>R</u> <u>W</u> | | | | |
| | | | Abundance: FC | FC | C |

STATUS

R-Resident (breeds)
N-Nonbreeder (resident)
B-Definite breeder
b-Likely breeder
n-Non-breeder (summer)
W-Winter visitor
M-Migrant

ABUNDANCE

| | |
|--------------------|--------------------|
| Ab - Abundant | R - Rare |
| C - Common | Und - Undetermined |
| FC - Fairly Common | Irr - Irregular |
| U - Unusual | |

| | | Breeding | Migr. | Winter | |
|---|--|---------------|----------|-----------|--------|
| | Habitat: | CF | R,CF,U | R,U,PP,PJ | |
| | Abundance: | FC | U-C | U-C | |
| Evening Grosbeak <u>Coccothraustes vespertinus</u> | M M M R b W — W R R R R — B B B R R W M R R b R B — W | | | | |
| Acc. in 21 | | | | | |
| House Sparrow <u>Passer domesticus</u> | R | Habitat: Ag,U | Breeding | Migr. | Winter |
| (Introduced) | | | | | |
| | | Abundance: Ab | | | Ab |

BIRD LATILONG - INDEX

Pg.

| | |
|--------------------------------|----|
| Anhinga..... | ii |
| Ani, Groove-billed..... | 31 |
| Avocet, American..... | 20 |
| Barn-owl, Common..... | 31 |
| Bittern, American..... | 3 |
| Least..... | 3 |
| Blackbird, Brewer's..... | 68 |
| Red-winged..... | 67 |
| Rusty..... | 68 |
| Yellow-headed..... | 68 |
| Black-hawk, Common..... | 13 |
| Bluebird, Eastern..... | 47 |
| Mountain..... | 47 |
| Western..... | 47 |
| Bobolink..... | 67 |
| Bobwhite, Northern..... | 17 |
| Brambling..... | 70 |
| Brant..... | 6 |
| Bufflehead..... | 10 |
| Bunting, Indigo..... | 61 |
| Lark..... | 64 |
| Lazuli..... | 61 |
| Painted..... | 61 |
| Snow..... | 67 |
| Bushtit..... | 44 |
| Canvasback..... | 8 |
| Cardinal, Northern..... | 60 |
| Catbird, Gray..... | 49 |
| Chat, Yellow-breasted..... | 59 |
| Chickadee, Black-capped..... | 44 |
| Mountain..... | 44 |
| Chukar..... | 15 |
| Coot, American..... | 18 |
| Cormorant, Double-crested..... | 2 |
| Olivaceous..... | 3 |
| Cowbird, Brown-headed..... | 69 |
| Crane, Sandhill..... | 18 |
| Whooping..... | 19 |
| Creeper, Brown..... | 45 |
| Crossbill, Red..... | 71 |
| White-winged..... | 71 |
| Crow, American..... | 43 |
| Cuckoo, Black-billed..... | 30 |
| Yellow-billed..... | 30 |
| Curlew, Eskimo..... | ii |
| Curlew, Long-billed..... | 21 |
| Dickcissel..... | 61 |
| Dipper, American..... | 46 |

| | |
|-------------------------------|----|
| Dove, Mourning..... | 30 |
| Rock..... | 30 |
| White-winged..... | 30 |
| Dowitcher, Long-billed..... | 24 |
| Short-billed..... | 24 |
| Duck, American Black..... | 7 |
| Black-bellied Whistling..... | 5 |
| Harlequin..... | 9 |
| Ring-necked..... | 9 |
| Ruddy..... | 11 |
| Wood..... | 7 |
| Dunlin..... | 24 |
| Eagle, Bald..... | 12 |
| Golden..... | 14 |
| Egret, Cattle..... | 4 |
| Great..... | 3 |
| Reddish..... | ii |
| Snowy..... | 4 |
| Falcon, Peregrine..... | 14 |
| Prairie..... | 15 |
| Finch, Cassin's..... | 70 |
| House..... | 70 |
| Purple..... | 70 |
| Rosy..... | 70 |
| Flicker, Northern..... | 37 |
| Flycatcher, Alder..... | ii |
| Ash-throated..... | 40 |
| Dusky..... | 38 |
| Dusky-capped..... | ii |
| Gray..... | 39 |
| Great Crested..... | 40 |
| Hammond's..... | 38 |
| Least..... | 38 |
| Olive-sided..... | 37 |
| Scissor-tailed..... | 40 |
| Vermilion..... | 39 |
| Western..... | 39 |
| Willow..... | 38 |
| Frigatebird, Magnificent..... | 3 |
| Gadwall..... | 8 |
| Gallinule, Purple..... | 18 |
| Gnatcatcher, Blue-gray..... | 47 |
| Godwit, Hudsonian..... | 22 |
| Marbled..... | 22 |
| Goldeneye, Barrow's..... | 10 |
| Common..... | 10 |
| Goldfinch, American..... | 71 |
| Lesser..... | 71 |
| Goose, Canada..... | 6 |
| Greater White-fronted..... | 6 |

| | <u>Pg.</u> |
|---------------------------------|------------|
| Ross'..... | 6 |
| Snow..... | 6 |
| Goshawk, Northern..... | 13 |
| Grackle, Common..... | 69 |
| Great-tailed..... | 68 |
| Grebe, Clark's..... | 2 |
| Grebe, Eared..... | 2 |
| Horned..... | 1 |
| Pied-billed..... | 1 |
| Red-necked..... | 2 |
| Western..... | 2 |
| Grosbeak, Black-headed..... | 61 |
| Blue..... | 61 |
| Evening..... | 71 |
| Pine..... | 70 |
| Rose-breasted..... | 60 |
| Grouse, Blue..... | 15 |
| Ruffed..... | 16 |
| Sage..... | 16 |
| Sharp-tailed..... | 16 |
| Gull, Bonaparte's..... | 26 |
| California..... | 27 |
| Franklin's..... | 26 |
| Glaucous..... | 28 |
| Glaucous-winged..... | 27 |
| Great Black-backed..... | 28 |
| Herring..... | 27 |
| Ivory..... | ii |
| Laughing..... | 26 |
| Lesser Black-backed..... | 27 |
| Little..... | 26 |
| Mew..... | 26 |
| Ring-billed..... | 27 |
| Ross'..... | 28 |
| Sabine's..... | 28 |
| Thayer's..... | 27 |
| Gyrfalcon..... | 15 |
| Harrier, Northern..... | 12 |
| Hawk, Broad-winged..... | 13 |
| Common, Black..... | 13 |
| Cooper's..... | 12 |
| Ferruginous..... | 14 |
| Red-shouldered..... | 13 |
| Red-tailed..... | 13 |
| Rough-legged..... | 14 |
| Sharp-shinned..... | 12 |
| Swainson's..... | 13 |
| Heron, Black-crowned Night..... | 4 |
| Great Blue..... | 3 |
| Green-backed..... | 4 |

| | |
|----------------------------------|----|
| Little Blue..... | 4 |
| Tricolored Heron..... | 4 |
| Yellow-crowned Night..... | 5 |
| Hummingbird, Anna's..... | 35 |
| Black-chinned..... | 35 |
| Blue-throated..... | 34 |
| Broad-tailed..... | 35 |
| Calliope..... | 35 |
| Magnificent..... | 34 |
| Rufous..... | 35 |
| Ibis, Glossy..... | 5 |
| White..... | 5 |
| White-faced..... | 5 |
| Jaeger, Long-tailed..... | ii |
| Parasitic..... | 26 |
| Pomarine..... | 25 |
| Jay, Blue..... | 42 |
| Gray..... | 42 |
| Pinyon..... | 43 |
| Scrub..... | 42 |
| Steller's..... | 42 |
| Junco, Dark-eyed..... | 66 |
| Kestrel, American..... | 14 |
| Killdeer..... | 20 |
| Kingbird, Cassin's..... | 40 |
| Eastern..... | 40 |
| Western..... | 40 |
| Kingfisher, Belted..... | 35 |
| Kinglet, Golden-crowned..... | 46 |
| Ruby-crowned..... | 47 |
| Kite, Mississippi..... | 12 |
| Swallow-tailed..... | 12 |
| Kittiwake, Black-legged..... | 28 |
| Knot, Red..... | 22 |
| Lark, Horned..... | 41 |
| Longspur, Chestnut-collared..... | 67 |
| Lapland..... | 67 |
| McCown's..... | 67 |
| Loon, Arctic..... | 1 |
| Common..... | 1 |
| Red-throated..... | 1 |
| Yellow-billed..... | 1 |
| Magpie, Black-billed..... | 43 |
| Mallard..... | 7 |
| Martin, Purple..... | 41 |
| Meadowlark, Eastern..... | 68 |
| Western..... | 68 |
| Merganser, Common..... | 11 |
| Hooded..... | 11 |
| Red-breasted..... | 11 |

| | <u>Pg.</u> |
|---------------------------------|------------|
| Merlin..... | 14 |
| Mockingbird, Northern..... | 49 |
| Moorhen, Common..... | 18 |
| Murrelet, Ancient..... | ii |
| Marbled..... | 29 |
| Nighthawk, Common..... | 32 |
| Lesser..... | 33 |
| Night-heron, Black-crowned..... | 4 |
| Yellow-crowned..... | 5 |
| Nutcracker, Clark's..... | 43 |
| Nuthatch, Pygmy..... | 45 |
| Red-breasted..... | 44 |
| White-breasted..... | 44 |
| Oldsquaw..... | 9 |
| Oriole, Northern..... | 69 |
| Orchard..... | 69 |
| Scott's..... | 69 |
| Osprey..... | 11 |
| Ovenbird..... | 57 |
| Owl, Barn..... | ii |
| Boreal..... | 33 |
| Burrowing..... | 32 |
| Common Barn..... | 31 |
| Eastern Screech..... | 31 |
| Flammulated..... | 31 |
| Great Horned..... | 32 |
| Long-eared..... | 32 |
| Northern Pygmy..... | 32 |
| Northern Saw-whet..... | 33 |
| Short-eared..... | 33 |
| Snowy..... | 32 |
| Spotted..... | 32 |
| Western Screech..... | 31 |
| Parula, Northern..... | 53 |
| Pewee, Eastern Wood..... | 38 |
| Western Wood..... | 38 |
| Pelican, American White..... | 2 |
| Brown..... | ii |
| Phainopepla..... | 50 |
| Phalarope, Red-necked..... | 25 |
| Red..... | 25 |
| Wilson's..... | 25 |
| Pheasant, Ring-necked..... | 15 |
| Phoebe, Black..... | 39 |
| Eastern..... | 39 |
| Say's..... | 39 |
| Pigeon, Band-tailed..... | 30 |
| Pintail, Northern..... | 7 |
| Pipit, Sprague's..... | 50 |
| Water..... | 50 |

| | |
|-------------------------------|----|
| Plover, Black-bellied..... | 19 |
| Lesser Golden..... | 19 |
| Mountain..... | 20 |
| Piping..... | 19 |
| Semipalmated..... | 19 |
| Snowy..... | 19 |
| Poorwill, Common..... | 33 |
| Prairie-Chicken, Greater..... | 16 |
| Lesser..... | 16 |
| Ptarmigan, White-tailed..... | 15 |
| Pygmy-owl, Northern..... | 15 |
| Quail, Gambel's..... | 17 |
| Scaled..... | 17 |
| Rail, Black..... | 17 |
| King..... | 17 |
| Virginia..... | 18 |
| Yellow..... | 17 |
| Raven, Chihuahuan..... | 43 |
| Common..... | 43 |
| Redhead..... | 9 |
| Redpoll, Common..... | 71 |
| Restart, American..... | 57 |
| Painted..... | 59 |
| Roadrunner, Greater..... | 31 |
| Robin, American..... | 49 |
| Ruff..... | 24 |
| Sanderling..... | 22 |
| Sandpiper, Baird's..... | 23 |
| Buff-breasted..... | 24 |
| Least..... | 23 |
| Pectoral..... | 23 |
| Semipalmated..... | 22 |
| Sharp-tailed..... | 23 |
| Solitary..... | 21 |
| Spotted..... | 21 |
| Stilt..... | 24 |
| Western..... | 23 |
| Upland..... | 21 |
| White-rumped..... | 23 |
| Sapsucker, Rednaped..... | 36 |
| Williamson's..... | 36 |
| Yellow-bellied..... | 36 |
| Scaup, Greater..... | 9 |
| Lesser..... | 9 |
| Scoter, Black..... | 10 |
| Surf..... | 10 |
| White-winged..... | 10 |
| Screech-owl..... | 31 |
| Western..... | 31 |
| Shoveler, Northern..... | 8 |

| | <u>Pg.</u> |
|-----------------------------|------------|
| Shrike, Loggerhead..... | 51 |
| Northern..... | 51 |
| Siskin, Pine..... | 71 |
| Snipe, Common..... | 25 |
| Solitaire, Townsend's..... | 47 |
| Sora..... | 18 |
| Sparrow, American Tree..... | 62 |
| Baird's..... | 64 |
| Black-throated..... | 64 |
| Brewer's..... | 63 |
| Cassin's..... | 62 |
| Chipping..... | 63 |
| Clay-colored..... | 63 |
| Field..... | 63 |
| Fox..... | 65 |
| Golden-crowned..... | 66 |
| Grasshopper..... | 64 |
| Harris'..... | 66 |
| Henslow's..... | 65 |
| House..... | 71 |
| Lark..... | 63 |
| Le Conte's..... | 65 |
| Lincoln's..... | 65 |
| Rufous-crowned..... | 62 |
| Sage..... | 64 |
| Savannah..... | 64 |
| Song..... | 65 |
| Sharp-tailed..... | 65 |
| Swamp..... | 66 |
| Vesper..... | 63 |
| White-crowned..... | 66 |
| White-throated..... | 66 |
| Spoonbill, Roseate..... | ii |
| Starling, European..... | 51 |
| Stilt, Black-necked..... | 20 |
| Stork, Wood..... | ii |
| Swallow, Bank..... | 41 |
| Barn..... | 42 |
| Cliff..... | 42 |
| Northern Rough-winged..... | 41 |
| Tree..... | 41 |
| Violet-green..... | 41 |
| Swan, Trumpeter..... | 6 |
| Tundra..... | 5 |
| Swift, Black..... | 34 |
| Chimney..... | 34 |
| White-throated..... | 34 |
| Tanager, Hepatic..... | 60 |
| Scarlet..... | 60 |
| Summer..... | 60 |
| Western..... | 60 |

| | Pg. |
|----------------------------|-----|
| Teal, Blue-winged..... | 7 |
| Cinnamon..... | 8 |
| Green-winged..... | 7 |
| Tern, Arctic..... | 29 |
| Black..... | 29 |
| Caspian..... | 28 |
| Common..... | 29 |
| Forster's..... | 29 |
| Least..... | 29 |
| Thrasher, Bendire's..... | 49 |
| Brown..... | 49 |
| Curve-billed..... | 50 |
| Long-billed..... | ii |
| Sage..... | 49 |
| Thrush, Gray-cheeked..... | 48 |
| Hermit..... | 48 |
| Swainson's..... | 48 |
| Varied..... | 49 |
| Wood..... | 48 |
| Titmouse, Plain..... | 44 |
| Towhee, Brown..... | 62 |
| Green-tailed..... | 62 |
| Rufous-sided..... | 62 |
| Turkey, Wild..... | 16 |
| Turnstone, Ruddy..... | 22 |
| Veery..... | 48 |
| Vireo, Bell's..... | 51 |
| Gray..... | 51 |
| Philadelphia..... | 52 |
| Red-eyed..... | 52 |
| Solitary..... | 52 |
| Warbling..... | 52 |
| White-eyed..... | 51 |
| Yellow-throated..... | 52 |
| Vulture, Turkey..... | 11 |
| Warbler, Bay-breasted..... | 56 |
| Black-and-white..... | 57 |
| Blackburnian..... | 55 |
| Blackpoll..... | 56 |
| Black-throated Blue..... | 54 |
| Black-throated Gray..... | 55 |
| Black-throated Green..... | 55 |
| Blue-winged..... | 52 |
| Canada..... | 59 |
| Cape May..... | 54 |
| Cerulean..... | ii |
| Chestnut-sided..... | 54 |
| Connecticut..... | 58 |
| Golden-winged..... | 53 |
| Grace's..... | 56 |

| | Pg. |
|-----------------------------|-----|
| Hermit..... | 55 |
| Hooded..... | 59 |
| Kentucky..... | 58 |
| Lucy's..... | ii |
| MacGillivray's..... | 58 |
| Magnolia..... | 54 |
| Mourning..... | 58 |
| Nashville..... | 53 |
| Orange-crowned..... | 53 |
| Palm..... | 56 |
| Paruva, Northern..... | 53 |
| Pine..... | 56 |
| Prairie..... | 56 |
| Prothonotary..... | 57 |
| Swainson's..... | 57 |
| Tennessee..... | 53 |
| Townsend's..... | 55 |
| Virginia's..... | 53 |
| Wilson's..... | 59 |
| Worm-eating..... | 57 |
| Yellow..... | 54 |
| Yellow-rumped..... | 54 |
| Yellow-throated..... | 55 |
| Waterthrush, Louisiana..... | 58 |
| Northern..... | 58 |
| Waxwing, Bohemian..... | 50 |
| Cedar..... | 50 |
| Whimbrel..... | 21 |
| Whip-poor-will..... | 34 |
| Wigeon, American..... | 8 |
| Eurasian..... | 8 |
| Willet..... | 21 |
| Woodcock, American..... | 25 |
| Woodpecker, Downy..... | 37 |
| Hairy..... | 37 |
| Ladder-backed..... | 37 |
| Lewis'..... | 36 |
| Red-bellied..... | 36 |
| Red-headed..... | 36 |
| Three-toed..... | 37 |
| Wren, Bewick's..... | 45 |
| Canyon..... | 45 |
| Carolina..... | 45 |
| House..... | 46 |
| Marsh..... | 46 |
| Rock..... | 45 |
| Sedge..... | 46 |
| Winter..... | 46 |
| Yellowlegs, Greater..... | 20 |
| Lesser..... | 20 |
| Yellowthroat, Common..... | 59 |

APPENDIX I

COLORADO LATILONG STUDY

CHANGE FORM

Species: _____

Please use this form to supplement or to correct data appearing in
The Colorado Bird Distribution Latilong Study (1982).

RECOMMENDED CODES

Latilong #: _____

Distribution Code: _____

Habitat Code: _____

Abundance: Abundant Common Fairly Com.

Unusual Rare Irregular

Undetermined

* * *

BASIS FOR RECOMMENDATION

Locality (Include nearest town & county) _____

Date(s) of observation _____

Number of birds _____

Description of observation; explain
justification for recommended code: _____

Name (print): _____

Mailing Address: _____

Send form to:

Zoology Dept.
Denver Museum of Natural History
2001 Colorado Blvd.
Denver CO 80205

Zip

APPENDIX II

COLORADO LATILONG STUDY

SCHEDULE OF RECOMMENDED CHANGES

Field worker

Address

Zip

COLORADO FIELD ORNITHOLOGISTS - SIGHT RECORD FORM

NEW STATE RECORD UNUSUAL RECORD

Please print or type. Attach photographs and additional pages if necessary.

Species: _____ / _____
(Vernacular Name) (Scientific Name)

Date(s): _____

Number of birds seen: _____ Sex: _____ Plumage: _____

Latilong: _____ Distribution Code: _____ Habitat Code: _____

Locality: _____ Elevation: _____

Nearest town: _____ County: _____

Time(s) observed: _____

Duration of observation (e.g. 1 sec., 5 min., etc.): _____

Other observers who saw the bird with you (give names, addresses, telephone numbers):

Other observers who saw the bird independently (give names, addresses, telephone numbers):

Light conditions (position of bird to light and shade, direction and strength of light):

Optical equipment used (e.g. power, types): _____

Distance and how measured: _____

If photographed, type of equipment and film: _____

Who else photographed the bird: _____

Describe the bird's behavior (e.g. flying, feeding, or habits used in identifying):

Describe in detail the bird's relative size, overall shape, plumage and color pattern. Include details of bill, wing and tail shape. Give particular emphasis to marks you used to identify the bird. Reference to its resemblance to field guide descriptions and illustrations is not enough. Describe only what you actually observed in the field:

Description (continued): _____

Describe the bird's song and call, if given, including method of delivery, (i.e. from perch, in flight, duration):

Describe the general and specific habitat in which you observed the bird:

How did you eliminate similarly appearing species?

What is your prior experience with this and similarly appearing species?

List books, illustration, recordings, other birders, etc. consulted and how this influenced your identification:

a) At time of observation: _____

b) After observation: _____

This report was written from reports made during _____, after observation _____, from memory _____

Date and time report was written: _____

Reporting observer: _____ Signature: _____

Address: _____

All reports will be maintained permanently at the address below:

Return to: CURATOR, DEPARTMENT OF ZOOLOGICAL COLLECTIONS
DENVER MUSEUM OF NATURAL HISTORY
CITY PARK
DENVER, CO 80205

RECEIVED

JUN 19 1989

COLORADO S.^tA.
State Publications Library