

HSR 18: Weld	Health Statistic Region: 18		Colorado (state estimate)				
	2013 % (weighted)	95% CI	2013 % (weighted)	95% CI			
Cigarette Use							
Ever-smoked	25.9	17.7-34.0	21.8	20.6-23.0			
Current smoker (overall)	13.4	8.4-18.3	10.7	10.0-11.4			
Sex							
Male	16.1	8.5-23.8	12.3	11.2-13.3			
Female	10.7	6.9-14.5	9.1	8.3-9.9			
Race/Ethnicity							
White (non-Hispanic)	12.3	5.2-19.3	10.1	9.2-10.9			
Hispanic/Latino	11.5	6.1-16.8	11.0	9.7-12.4			
African American / Black		•	10.6	7.0-14.2			
Asian American		•	6.9	4.9-8.8			
Native Hawaiian or Other Pacific Islander	·	•	10.9	4.8-17.0			
American Indian/Alaska Native			22.8	17.3-28.4			
Other	37.0	26.2-47.8	17.9	15.3-20.5			

Frequent smoker (smoked 20-30 days in past 30)	5.4	4.6-6.2	3.3	2.9-3.7		
Quit attempt in the past year (among smokers)	42.0	32.5-51.6	50.8	47.4-54.3		
Probably or definitely smoke in the next year	16.9	8.8-25.0	14.1	13.3-14.9		
Non-cigarette tobacco use						
Current smokeless user	10.9	9.4-12.3	6.4	5.9-6.9		
Current cigar user	10.4	7.2-13.5	9.7	8.9-10.6		
Ever tried e-cigarettes	13.8	6.1-21.5	15.1	14.1-16.1		
Never tried other tobacco products	45.4	39.0-51.8	46.8	45.2-48.4		
Secondhand smoke exposure						
Secondhand smoke, room (past 7 days)	16.7	12.0-21.5	15.5	14.4-16.6		
Secondhand smoke, car, parent (past 7 days)	13.5	8.9-18.1	13.2	12.1-14.3		
Think secondhand smoke is moderate/great risk	80.6	77.8-83.4	76.2	75.1-77.3		

. = Less than 50 students in the subgroup.

